


UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 5605-110



CHIGGERS AND CERTAIN LAWN INSECTS: Spray lawns, picnic areas and infested foliage sufficiently to cover area.

CLOVER MITES: Spray window and door sills, foundation and areas around foundation.

CAUTION: Harmful if swallowed. May be absorbed through the skin. Avoid inhalation of spray mist and skin contact. In case of contact, wash immediately with soap and water. For eyes, flush with water for 15 minutes. Get medical attention. Do not use on household pets or humans. Avoid contamination of feed and food-stuffs. Keep out of reach of children. Do not use or store near heat or open flame.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment, or disposal of wastes.

ACTIVE INGREDIENTS:	By Weight
*Malathion	19.5%
**Methoxychlor, Technical	15.0%
***1,1-bis (p-chlorophenyl)-2,2,2-trichloroethanol	0.9%
Methylated Naphthalenes	59.6%
INERT INGREDIENTS	5.0%
	Total—100.0%

- *O,O-dimethyl Dithiophosphate of diethyl mercaptosuccinate.
- **Equivalent to 13.2% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 1.8% related compounds.
- ***Kelthane, registered trademark of Rohm & Haas.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

NET CONTENTS 1 LIQUID QUART

257

MANUFACTURED FOR
FARMINGDALE GARDEN LABS. INC.
136 VERDE STREET, FARMINGDALE, N.Y. 11735

U.S. PAT. REG. NO. 160,000

NET CONTENTS 1 LIQUID PINT

256

HOME AND GARDEN INSECT SPRAY

USE ON: Roses, Flowers, Vegetable Crops, Fruit Trees, Shrubs, Evergreens and Lawns

KILLS: Aphids, Armyworms, Azalea Scale, Birch Leaf Miners, Black Scale Crawlers, Bud Moths, Chiggers, Codling Moths, Clover Mites, Cucumber Beetles, Elm Leaf Miners, Euonymus Scale, Flea Beetles, Japanese Beetles, Leafhoppers, Mealy Bugs, Mexican Bean Beetles, Oriental Fruit Moths, Oyster Shell Scale, Plant Bugs, Plum Curculio, Red Spiders, Soft Brown Scale, Spider Mites, Spittle Bugs, Tent Caterpillars, Thrips, White Flies.

CAUTION: KEEP OUT OF REACH OF CHILDREN.

See other warnings on side panel

Cont

FARMINGDALE
Malathion
formula
insects
leaves
growing
interval
gallon

FLOWER
flowering
oli, ev
shrubs
leafhoppers
and el
crawlers
bugs, c
worms,

VEGETABLE
beetles
hopper

FRUIT
plums,
rotten,
Japanese
and so
mended
7 days
21 days
and Ca
may b

FLIES
porches
where v
stagnant

4 21 10
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 5605-110



FARMINGDALE GARDEN LABORATORIES, INC.
Farmingdale

PROVEN PRODUCT

HOME AND GARDEN INSECT SPRAY

**USE ON: Roses, Flowers, Vegetable Crops,
Fruit Trees, Shrubs, Evergreens and Lawns**

**KILLS: Aphids, Armyworms, Azalea Scale, Birch Leaf Miners,
Black Scale Crawlers, Bud Moths, Chiggers, Codling Moths,
Clover Mites, Cucumber Beetles, Elm Leaf Miners, Euonymus
Scale, Flea Beetles, Japanese Beetles, Leafhoppers, Mealy Bugs,
Mexican Bean Beetles, Oriental Fruit Moths, Oyster Shell Scale,
Plant Bugs, Plum Curculio, Red Spiders, Soft Brown Scale, Spider
Mites, Spittle Bugs, Tent Caterpillars, Thrips, White Flies.**

CAUTION: KEEP OUT OF REACH OF CHILDREN.

See other warnings on side panel

Contains Malathion, Kelthane and Methoxychlor

FARMINGDALE HOME AND GARDEN INSECT SPRAY, which contains Malathion, Methoxychlor and Kelthane, is a specially prepared formulation for the control of fruit, vegetable and ornamental insects and mites. Apply to both upper and underneath sides of leaves until plants are dripping wet. Protect plants during the growing season by repeating spray applications at 7 to 14 day intervals or as necessary. Dilute 1 oz. (2 tablespoonsful) to one gallon of water for spray mixture.

FLOWERS, ORNAMENTALS, SHRUBS AND TREES: Spray roses, flowering plants, chrysanthemums, dahlias, camellias, asters, gladioli, evergreens, azaleas, rhododendrons, spirea, lilacs, and other shrubs for control of Japanese beetles, aphids, spider mites, thrips, leafhoppers, flea beetles, mealy bugs, spittle bugs, birch, boxwood and elm leaf miners, plant bugs, soft brown scale, black scale, crawlers, oyster shell scale, euonymus scale, azalea scale, lace bugs, armyworms, tent caterpillars, leaf roller caterpillars, cankerworms, red spiders, white flies, and certain other insects.

VEGETABLES: Use on tomatoes, peppers, cucumbers, beans, for flea beetles, aphids, Mexican bean beetles, cucumber beetles, leafhoppers. Do not use within 7 days of harvest.

FRUIT TREES: Spray apples, apricots, peaches, pears, cherries, plums, and grapes for protection against aphids, red-banded leaf roller, codling moth, spider mites, plum curculio leafhoppers, Japanese beetles, tent caterpillars, oriental fruit moths, bud moths and soft brown scale. Apply at 7 to 10 day intervals or as recommended by State Extension Service Specialists. Do not use within 7 days of harvest. For apricots and peaches, do not use within 21 days of harvest; grapes within 14 days of harvest. MacIntosh and Cortland apples, certain varieties of sweet cherries, Bosc pears may be injured.

FLIES AND MOSQUITOES: Spray garbage cans, indoor drains, porches and foliage around patio areas. Mosquito breeding areas, where water stands, use this dosage only in intermittently flooded areas, stagnant water and temporary rain pools.