

culio, Leafhoppers, Japanese Beetle Adults, Oriental Fruit Moths, Bud Moths: 1 tablespoonful per gallon of water. Scales: 1½ to 2 tablespoonfuls per gallon of water. Apply at 7 to 10 day intervals or as recommended by State Agricultural Experiment Station. Do not use within 7 days of harvest on peaches & apricots. This product may injure McIntosh and Cortland varieties of apples if applied within 4 weeks of harvest.

FLIES and MOSQUITOS: 5 tablespoonfuls per gallon of water. Spray patios, foliage and grass around patios and porches at the rate of 1 gallon per 1000 sq. ft.

CLOVER MITES, CHIGGERS, CRICKETS, EARWIGS and certain **LAWN INSECTS:** 8 tablespoonfuls per gal. of water. Spray foundations, window and door sills, lawn and picnic areas at the rate of 1 gal. of dilute spray per 100 sq. ft.

CAUTION: Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin; wash thoroughly after using. Avoid contamination of feed and foodstuffs.

KEEP OUT OF REACH OF CHILDREN.

Highly toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate ornamental fish ponds.

Apply this product only as specified on this label.

FARMINGDALE GARDEN LABS. INC.
136 Verdi Street, Farmingdale, L. I., N.Y. 11735
U.S.P.A. REG. NO. 5605-115

KILLS EARWIGS



FARMINGDALE GARDEN LABORATORIES, INC.

Farmingdale

PROVEN PRODUCT

NEW IMPROVED

MALATHION 50%

KILLS: Insects, Mites & Scales.

USE ON: Roses, Flowers, Evergreens, Shrubs, Trees, Fruits, and Vegetables.

CAUTION:

KEEP OUT OF REACH OF CHILDREN

See other caution on side panel

NET CONTENTS 1 LIQUID QUART

ACTIVE INGREDIENTS:

Malathion*	By Weight	50%
Aromatic Petroleum Derivative Solvent		45%

INERT INGREDIENTS	5%
-------------------	----

TOTAL 100%

*O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

MAKES
UP TO
128
GALLONS
MIXED
SPRAY

For Use in Hose Sprayer, Tank Sprayer or Watering Can.
2 tablespoonfuls equals 1 ounce.
3 teaspoonfuls equals 1 tablespoonful.

DIRECTIONS: Apply to both upper and lower leaf surfaces and other plant parts. Repeat spraying at 7 to 14 day intervals, or as necessary. Do not apply to food crops within 3 days of harvest or as otherwise specified.

ORNAMENTALS: (Roses, Flowers, Evergreens, Shrubs, and Trees). Aphids, 1½ to 2 teaspoonfuls per gallon of water. Mites, Red Spider Mites, Spruce Mites, Japanese Beetle Adults, Mealy Bugs, Thrips, Leafhoppers, Flea Beetles, White Fly, Lace Bug, Bagworm, Tarnished Plant Bug, Birch, Boxwood and Elm Leaf Miners, Tent Caterpillar: 1 tablespoonful per gallon of water. Soft Brown Scale, Black Scale Crawlers, Oyster Shell Scale, Euonymus Scale, Azalea Scale: 1½ to 2 tablespoonfuls per gal. of water. Injury may occur to certain ferns including Boston, Maidenhair and Pteris as well as some species of Crassula.

VEGETABLES: (Tomatoes, Peppers, Cucumbers, Beans, Cabbage and Cauliflower). Flea Beetles, Aphids, Mexican Bean Beetles, Cucumber Beetles, Leafhoppers, Red Spider Mites, Thrips, Cabbage Worms: 1 tablespoonful per gallon of water. Do not use within 7 days of harvest on cabbage and Cauliflower.

FRUIT TREES and BUSH FRUITS (Apples, Pears, Peaches, Apricots, Cherries, Grapes and Blackberries): Aphids, Red-Banded Leaf Rolier, Codling Moth, Mites, Red Spider Mites, Plum Cur

ACCEPTED

NOV-18-1970

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE ACT
FOR THE PROTECTION OF
UNDER 5605-115