




CAUTION: DO NOT USE, POUR, SPILL OR STORE
NEAR HEAT OR OPEN FLAME.

This concentrate, if brought into contact with the skin can cause toxic symptoms. Avoid skin contact and inhalation. In case of spillage on skin, wash immediately with soap and water. Avoid contamination of feed, food containers and foodstuffs. Do not use on household pets or humans. HARMFUL IF SWALLOWED. Keep out of reach of children. Do not use this product on dairy animals, in dairy barns, or on feed or forage to be fed to dairy animals or other animals being finished for slaughter. Do not use in poultry houses. Toxic to fish and wildlife. Birds feeding on treated area may be killed. Do not contaminate any body of water nor apply when conditions favor runoff from area treated. Do not contaminate water by cleaning of equipment or disposal of waste. Do not reuse container when empty. Destroy by burying.



CHLORDANE TERMITE PROOFER

ACTIVE INGREDIENTS:
Chlordane - Technical* 45.0%
Kerosene 50.0%
INERT INGREDIENTS 5.0%
Total.....100.0%
(Contains 4 lbs. Chlordane-Tech per U.S. gal. at 70°F.)
*27% Octachloro-4,7-methanotetrahydroindane and
18% related compounds.

CAUTION: Keep Out of Reach of Children

SEE REAR PANEL FOR ADDITIONAL CAUTIONS

NET CONTENTS 1 QUART FLUID

E.P.A. REG. No. 5605-140
E.P.A. EST. No. 5605-N.Y.-2

110A

Farmingdale Garden Labs, Inc.

136 VERDI STREET
FARMINGDALE, L.I., N.Y. 11735

SUBTERRANEAN TERMITES

DIRECTIONS FOR NON-PROFESSIONALS

Use FARMINGDALE CHLORDANE TERMITE PROOFER in a
Prepare this emulsion as follows: Use one quart of FARMINGDALE
TERMITE PROOFER in 12.5 gallons of water. Mix thoroughly.

BUILDINGS HAVING BASEMENTS OR CRAWL SPACES:

Dig a trench along the outside of the foundation walls, block or concrete foundations, dig a trench to, BUT NOT DEEPER THAN, the bottom of the foundation. Then as the trench is refilled, treat the soil at the rate of 4 lbs. per 10 linear feet FOR EACH FOOT OF DEPTH. A trench 3 feet deep will treat 10 linear feet.

The objective is to establish a treated soil barrier which will prevent termites from entering. Other critical areas may also require treatment. If in doubt, consult the State Extension Service for advice.

Apply this product only as specified on this label.



CHLORDANE TERMITE PROOFER

DO NOT USE, POUR, SPILL OR STORE
NEAR HEAT OR OPEN FLAME.

Direct contact with the skin can cause
irritation and inhalation. In case of
contact immediately with soap and water. Avoid
containers and foodstuffs. Do not
use on humans. HARMFUL IF SWALLOWED. Keep
from use this product on dairy ani-
mals. Do not feed or forage to be fed to dairy
animals being finished for slaughter.

Birds feeding on treated area may
be contaminated. Do not apply when
area treated.

by cleaning of equipment or disposal
container when empty. Destroy by burying.

ACTIVE INGREDIENTS:

Chlordane - Technical*	45.0%
Kerosene	50.0%
INERT INGREDIENTS	5.0%
Total	100.0%

(Contains 4 lbs. Chlordane-Tech per U.S. gal. at 70°F.)
*27% Octachloro-4,7-methanotetrahydroindane and
18% related compounds.

CAUTION: Keep Out of Reach of Children

SEE REAR PANEL FOR ADDITIONAL CAUTIONS

NET CONTENTS 1 QUART FLUID

E.P.A. REG. No. 5605-140
E.P.A. EST. No. 5605-N.Y.-2

110A

Farmingdale Garden Labs, Inc.

136 VERDI STREET
FARMINGDALE, L.I., N.Y. 11735

SUBTERRANEAN TERMITES

DIRECTIONS FOR NON-PROFESSIONAL USE

Use FARMINGDALE CHLORDANE TERMITE PROOFER in a 1% water emulsion.
Prepare this emulsion as follows: Use one quart of FARMINGDALE CHLORDANE
TERMITE PROOFER in 12.5 gallons of water. Mix thoroughly.

BUILDINGS HAVING BASEMENTS OR CRAWL SPACES:

Dig a trench along the outside of the foundation walls. In brick or hollow
block or concrete foundations, dig a trench to, BUT NOT BELOW, the footing.
Then as the trench is refilled, treat the soil at the rate of 4 gallons per 10 linear
feet FOR EACH FOOT OF DEPTH. A trench 3 feet deep would require 12 gal-
lons per 10 linear feet.

The objective is to establish a treated soil barrier which will prevent termite
entry. Other critical areas may also require treatment. It is suggested that you
consult the State Extension Service for advise.

Apply this product only as specified on this label.

TESTED

1976

Insecticide,
Fertilizer Act,
pesticide

5-140

IMPORTANT — PLEASE NOTE

Farmingdale A.M.A. plus 2,4-D selectively kills crabgrass and also kills most common lawn weeds such as chickweed, dandelion, plantain, knotweed, pursley, spotted spurge, large leaf spurge, pennywort, beggar weed and certain others. May reduce clover. For use on Bermuda, Merion Blue, Kentucky Blue, Red Top, Fescue, Zoysia. Bent lawns only if well watered prior to application and average day temperatures under 85°. Do not use on St. Augustine, Carpet, Centipede, Bahia grasses, bent putting greens or Dichondra lawns.

HOW TO CONTROL: Good soil moisture required for best results. If dry, water 1-24 hours before applying. Resume keeping soil moist day after application and especially during dry, hot weather if possible. Some temporary browning may develop under drought conditions. During drought, late afternoon applications advisable. Apply when grass leaves are dry. Always repeat at 5-7 day intervals. Continue until plants are dead. This may range from 1-2 treatments for small plants, up to 3-4 for established plants. Plants will be dead 4-5 days after last required treatment.

TURF APPEARANCE: If correct rate and coverage was employed, crabgrass should start to lighten in color within 3 days and some browning occur at edge of leaf. If not, check rate used and coverage per gallon obtained and make needed correction on subsequent applications. Severe browning of all turf indicates excess rate was used. Continue applications of lighter rates.

NEW LAWN: Do not treat until after third mowing.

HOW TO APPLY

- With Nozzle Type Sprayer which attaches to garden hose: First measure lawn area to be treated. Place recommended (see above) amount of A.M.A. plus 2,4-D in glass jar, then fill jar with water and spray uniformly over the area to be treated.
- With Pressure Type Sprayer (type carried on back): Mix recommended amount of A.M.A. plus 2,4-D as required (see above) with 3 gallons of water and spray uniformly over the area to be treated.

NET CONTENTS 32 OZ.

192

A.M.A. plus 2,4-D CRABGRASS AND LAWN WEED KILLER

ACTIVE INGREDIENTS:

Dodecyl Ammonium Methyl Arsonate	8.00%
Octyl Ammonium Methyl Arsonate	8.00%
*Octyl Ammonium Salt of 2,4-Dichlorophenoxyacetic acid..	5.44%
INERT INGREDIENTS	78.56%
Total	100.00%
Total arsenic, expressed as metallic, all in water soluble form	4.00%
*Equiv. to 2,4-Dichlorophenoxyacetic acid	3.43%

Manufactured for
FARMINGDALE GARDEN LABS., INC.
153 Florida St., Farmingdale, N. Y.

KEEP OUT OF REACH OF CHILDREN
Read All Directions and Cautions Before Use

DIRECTIONS

If temperature is over 85°F. when applications are made, make sure of soil moisture and use the low dosage of 16 oz. (one pint) diluted in 10 to 15 gallons water and apply to 2500 sq. ft. of lawn area. Follow with two more applications at 7 day interval. At lower temperatures apply 16 oz. for each 2000 sq. ft. for young crabgrass and the same amount for 1500 sq. ft. of established crabgrass. Be sure soil is moist before treating. During a drought, water grass well before applications are made. Keep lawn mowed at one inch to one and one-half inches during treatment. Normal rainfall after applications will not affect results. **IMPORTANT:** Measure the area to be treated before applications are made. Determine square footage to be treated by multiplying length by the width (for example, an area 20 feet wide by 50 feet long contains 1000 square feet).

WARNING

Do not allow spray to contact or drift into trees, shrubs, flowers or vegetable plants. Coarse sprays are less likely to drift. Do not store near fertilizers, seeds, insecticides or fungicides. Avoid prolonged breathing of dust or spray mist. Avoid contact with skin, eyes or clothing. Avoid contamination of feed and food-stuffs. Keep away from children and domestic animals. Keep livestock off treated areas. Use on grass only. Harmful if swallowed. Do not contaminate waters used for domestic or irrigational purposes.

ANTIDOTE: If taken internally, induce vomiting and call physician at once.

DISCLAIMER: Above representations are based on tests designed to protect turf from improper use. However, no guarantees expressed or implied, are made and buyer assumes all risk.

Product Patent applied for. Licensed for use in the control of crabgrass according to the method of U.S. Pat. No. 2,678,265 by reason of inclusion in the purchase price of a royalty. A license at the same royalty rate to use the patented method with unpatented material obtained from any source will be granted upon request by the patent owner.

Camerman's Note
Poor Copy

ACCEPTED

UNDER THE FEDERAL NARCOTICS
-UNIQUE AND TRUSTED-
FOR DOCTORS, PHARMACISTS AND
UNDER NO. 100-100000