


FRONT PANEL

ETCH

Technical Maintenance Products, Inc. - Philadelphia, Pa.

Net Content
1.0 liter (33.8 fl. oz.)
1.0 liter (33.8 fl. oz.)
1.0 liter (33.8 fl. oz.)
1.0 liter (33.8 fl. oz.)

Net Content
1.0 liter (33.8 fl. oz.)
1.0 liter (33.8 fl. oz.)



Keep out of reach of children.
Use only panel for other instructions.

Net Content

TECHNICAL MAINTENANCE PRODUCTS, INC.
2011 N. American St.
Phila., Pa. 19133

STERE PANEL

DIRECTIONS

Milking machinery, tanks and pails should be pre-cleaned before wetting thoroughly with an ETCH solution containing 25-50 and 50-100 ppm of titratable iodine. Allow solution to remain in contact with all surfaces to be disinfected for at least 10 minutes, then rinse all treated food contact surfaces with potable water prior to reuse.

Surgical Instruments - All blood and other adhering soil should be removed before disinfecting with an ETCH solution containing 100 to 150 ppm of titratable iodine. Immerse instruments for at least ten (10) minutes.

DILUTION CHART
(per 10 gallon water)

PPM TITRATABLE IODINE	VELOCITY OF ACTION
25	1
50	2
75	3
100	4
125	5
150	6

For hospital use on lean walls, floors, and equipment, use 100 ppm on most food surfaces and 150 ppm on porous surfaces.

CAUTION: This is an acid material. Do not use internally.
If swallowed, drink much milk or water and call physician.
Avoid contamination of food. May cause skin irritation. Avoid contact with skin or eyes. Flush eyes freely with water. Call physician.