

FOR INDUSTRIAL AND INSTITUTIONAL USE ONLY


TECH-CIDE

ACTIVE INGREDIENTS:

Petroleum distillates 84%
o-Isopropoxyphenyl
methylcarbamate* 1%

INERT INGREDIENTS: 15%

* Contains BAYGON, registered trademark of Farbenfabriken Bayer A.G.; Chemagro Corporation licensee. U.S. Patent No. 3,111,539. This material used under license from Chemagro Corporation.



CAUTION: Keep out of reach of children

May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing of spray mist and provide adequate ventilation of area being treated. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using. Avoid contamination of food, foodstuffs, food utensils, and food preparation areas. Do not allow children to contact treated areas until surfaces are thoroughly dry. Should poisoning occur, get prompt medical aid. To PHYSICIAN: Atropine Sulfate is antidotal.

NET CONTENTS:

GALLON

MANUFACTURED BY

TECHNICAL MAINTENANCE PRODUCTS, INC.
2211 N. American Street, Philadelphia, Pa. 19133



TECH-CIDE is a highly effective ready-to-use formulation for control of insects listed, including those having resistance to certain chlorinated hydrocarbons and phosphate insecticides. It possesses rapid kill, good flushing action, and prolonged residual activity. **KILLS:** ants, brown dog ticks, centipedes, cockroaches, clover mites, crickets, earwigs, firebrats, fleas, flies, gnats, millipedes, mosquitoes, paperlice, punkies, sandflies, scorpions, silverfish, sowbugs and spiders. Exposed stages of cadelles, cigarette beetles, dermestids, drugstore beetles, Indian meal moths, Mediterranean flour moths, red flour beetles, saw-toothed grain beetles, and weevils.

1

ACCEPTED

Mar. 23, 1971

UNDER THE FEDERAL INSECTICIDE ACT
 FOR LISTED PESTICIDE REGISTERED
 UNDER THE SUBJECT TO ATTACHED COMMENTS.

DIRECTIONS FOR USE

GENERAL: Apply TECH-CIDE at the rate of one (1) gallon per 1,000 sq. ft. Use a pressurized sprayer with a fine flat spray pattern. Coat surface with a dew-like mist. Apply TECH-CIDE to cracks, crevices, corners, perimeters of rooms, behind baseboards, around window and door frames; behind escutcheon plates; under stoves and sinks; to the under-surface of shelves, drawers and work tables; to elevator pits and shafts; behind wooden or metal panels, at seams, joints, and contiguous surfaces of equipment such as dishwashers and refrigerators; around electric motors and compressors; laundry carts; garbage areas and cans; warm dark moist areas; void spaces; hidden surfaces and quiet undisturbed areas. Apply TECH-CIDE to all harborages and breeding areas of insects, or to surfaces where they congregate or crawl. Repeat treatment as necessary to prevent or control infestation. Remove pets and cover fish bowls before spraying. Do not spray animals.

ANTS: Give special attention to doors, windows, and ant trails.

COCKROACHES: Thoroughly treat the infested area with special attention to cracks, crevices, around sinks, under surfaces and other hard-to-get-at places, as well as in food handling and storage areas and the harborages listed above.

CLOVER MITES, CRICKETS, EARWIGS, MILLIPEDES, CENTIPEDES, SOWBUGS, SCORPIONS, AND SPIDERS: Thoroughly spray the interior and exterior perimeter of the infested premises. For clover mites, give special attention to window frames. Do not treat foliage.

FLEAS, FLIES, GNATS, MOSQUITOES, PUNKIES, SANDFLIES (outdoors only): Thoroughly treat all outdoor surfaces where these insects are known to alight or congregate. Treat outside surfaces such as doors, window screens and around other areas where they may enter the building. Thorough coverage is necessary for maximum control.

BROWN DOG TICKS: Thoroughly treat floor areas, animal sleeping quarters, and wherever infestation is suspected. Provide fresh bedding. Do not spray animals. Have animals treated by a veterinarian.

ACCESSIBLE OR EXPOSED STAGES OF CADELLES, CIGARETTE BEETLES, DRUGSTORE BEETLES, INDIAN MEAL MOTHS, MEDITERRANEAN FLOUR MOTHS, RED FLOUR BEETLES, DERMESTIDS, SAW-TOOTHED GRAIN BEETLES AND WEEVILS: Thoroughly treat walls and floors of infested area with attention to cracks, crevices, corners, and perimeters. Sanitation and dust control are of primary importance. Fumigate or destroy infested merchandise.

SILVERFISH, FIREBRATS, AND PAPERLICE: Thoroughly treat baseboards, bookcases, and storage areas.

FOR PEST CONTROL IN MEAT AND POULTRY PROCESSING PLANTS: Use TECH-CIDE only in the inedible products areas of meat packing and food processing plants. Spraying loading docks and outside areas where insects congregate will aid in preventing reinfestation. Good sanitation and proper waste disposal are essential in any effective insect control program. Do not contaminate water by cleaning of equipment or disposal of wastes.

IN FOOD PROCESSING AND FOOD STORAGE AREAS: Treat food processing plants early in the season before canning and other processing start. Repeat treatment as necessary, when plant is not in operation. Spraying pallets, loading docks, and outside areas where insects congregate will aid in preventing reinfestation. Do not use in commercial food preparation areas or in edible products areas of food processing plants.

INSTITUTIONAL AND INDUSTRIAL: TECH-CIDE should be used around hospitals, (when patients are out of area), hotels, schools, theaters, housing developments, warehouses, transportation equipment such as trucks, ships, and railroad cars (when they are empty of produce or merchandise). Remove animals from the area when making applications around zoos.