

ALWAYS DILUTE ACCORDING TO
 "DILUTION TABLE" BEFORE USING
 DIRECTIONS FOR USE
 DO NOT ADD TO SOAP OR
 DETERGENT SOLUTION

1. For routine use. Swab, wipe, or rinse surfaces (such as floors, walls, urinals, washstands, garbage pails), which have been previously cleaned with soap or detergent solution and rinsed thoroughly, using one percent solution (see "DILUTION TABLE").
2. For special use. If known heavy contamination exists, use two percent solution on surfaces. Swab, wipe or rinse surfaces which have been previously cleaned with a soap or detergent solution and rinsed, with disinfectant solution at least twice, allowing solution to act for at least ten minutes between applications. Disinfect smaller objects by immersion for at least ten minutes. Permit all surfaces to dry before use.
3. Precaution. To disinfect RUBBER and PLASTIC objects, swab surfaces if possible. Immerse only if required. LIMIT EXPOSURE TO TEN MINUTES PROLONGED IMMERSION IN DISINFECTANT MAY AFFECT PLASTICS AND RUBBER SERIOUSLY
4. Rinse empty container thoroughly with water and discard it.

**DISINFECTANT, GERMICIDAL,
 CONCENTRATE
 FOR USE IN HOSPITALS,
 SCHOOLS, INDUSTRY &
 INSTITUTIONS**

ACTIVE INGREDIENTS:

Ortho-Benzyl-Para-Chlorophenol Sodium Salt 4.00%
 Sodium Dodecylbenzenesulfonate 3.00%
 Isopropanol 10.00%
 Essential Oils 0.25%

INERT INGREDIENTS: 82.75%

WARNING:

KEEP OUT OF REACH OF CHILDREN.
 DO NOT TAKE INTERNALLY, HARMFUL IF SWALLOWED.
 RAPIDLY ABSORBED THROUGH SKIN
 AVOID CONTACT WITH SKIN:
 WASH THOROUGHLY AFTER USING.
 CAUSES SEVERE BURNS.
 DO NOT GET IN EYES; ON SKIN, OR ON CLOTHING.
 AVOID BREATHING VAPOR.
 ALWAYS DILUTE ACCORDING TO "DILUTION TABLE" BEFORE USING.
 SEE FIRST AID ON PANEL SIDE.

EPA NO 5667-67
 6840-00-530-7109
 DISINFECTANT - TYPE I
 O-D-001435
 1-U. S. GALLON
 BARRETT CHEMICAL, CO., INC.
 H & LUZERNE STS. - PHILA., PA. 19124

FIRST AID

In case of contact with skin, wash thoroughly with soap and water; for eyes, flush with plenty of water for at least 15 minutes and get medical attention. If swallowed, administer milk or raw egg and give large amounts of water. Then cause vomiting by inserting finger down throat. See a medical doctor at once.

DILUTION TABLE

Dilution	Ounce of concentrate per gallon of water	Ml. concentrate per liter of water	Teaspoons concentrate per gallon of water
1 percent solution (Routine use)	1 1/4	10	10
2 percent solution (Special use)	2 1/2	20	20

ACCEPTED WITH COMMENTS