

TECHNICAL MAINTENANCE PRODUCTS, INC.

SIDE PANEL



SURE

WEED KILLER

DIRECTIONS (18 point)

Areas of application must be limited to bare ground or small enclosed industrial areas. For treating parking places, gutters, drives, walks, or other places where no growth is wanted dilute SURE WEED KILLER 1 - 40 parts of water. Apply carefully with sprinkling can or a similar spraying device; make sure all weeds are completely drenched. If ground is dry use water hose to thoroughly wet down area before applying SURE WEED KILLER. It is best to apply SURE WEED KILLER after it has rained. To kill thistles, crabgrass, poison ivy, or any other hard weeds, dilute SURE WEED KILLER 1 - 20 parts of water and spray thoroughly. If regrowth appears, a second application can be applied if desired.

Dosage rates in excess of 50 pounds of arsenic trioxide equivalent per acre are limited to areas to be covered by pavement after treatment.

CAUTION (18 point)

Keep off of treated areas, or any part of the skin. Keep cats, dogs, chickens, cattle, or any other grazing animal off treated areas for a period of three weeks or until a drenching rain has fallen on the sprayed area. Public roads should be posted with warning signs obtained from H&M. SURE WEED KILLER harms all vegetation; therefore, do not use on ornamentals, gardens, lawns, etc., unless you wish to kill them.

Branched Stage. Use 3 tablespoonfuls in 1 gallon water, apply evenly to 150 sq. ft. Make the second application 7 days later. In some cases additional applications may be necessary. Apply at 7 day intervals.

CAUTION: Harmful if swallowed. Avoid breathing spray mist. Avoid contact with eyes, skin and clothing. Wash thoroughly with soap and water after handling. Avoid contamination of feed and food stuff. Keep in original containers. Do not reuse empty container. When container is empty, wash thoroughly and destroy. Do not contaminate lakes, ponds, streams, woodlands and other non-crop areas.

If taken internally call a physician. Induce vomiting by giving patient a tablespoon of salt in a glass of warm water. Repeat until fluid is clear.


NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. **EPA REG. NO. 5719-30**

CHACON CHEMICAL CORPORATION
5245 Chakemco Street • South Gate, California 90280
A Subsidiary of Leisure Enterprises, Inc.

CHACON

CRABGRASS CONTROL

**For Use In:
DICHOANDA &
GRASS LAWNS**



ACTIVE INGREDIENTS	
Disodium Methearsonate	12.6%
INERT INGREDIENTS	
	87.4%

Total arsenic, all in water soluble form, expressed as elemental 5.12%
*Equivalent to 20% Disodium Methearsonate Hexahydrate.

CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions.

NET 16 FL. OZ. (1 PINT)

ACCEPTED

FEB 11 1974

UNDER THE FEDERAL INSECTICIDE AND ROENTGENIC ACT

REGISTERED UNDER EPA REG. NO. 5719-30

TO APPLICATOR (M.V.)

CHACON CRABGRASS CONTROL

Read label thoroughly. Use only in accordance with label cautions and directions.

Chacon Crabgrass Control is a post-emergence control; — kills both hairy and smooth-leaf Crabgrass. Crabgrass seeds start germinating in the spring about March or April. Plants mature through the summer and, being annuals, start turning brown and dying in late summer or early fall.

DIRECTIONS

For best results observe the following: Make first application as soon as possible after seeds start to germinate. Do not apply on grass or dichondra lawns under 2 months old. Water thoroughly before treatment and withhold water for 48 hours thereafter. Do not use on St. Augustine or centipede grasses.

How to Apply in the Seedling Stage (smaller than 2 inches). Use 2 tablespoonfuls in 1 gallon water, apply evenly to 150 sq. ft. Make the second application 7 days later. Uniform coverage is most essential.