

Citrus to control Red Scale, Citricola Scale, Brown Soft Scale. Start spraying in spring. Do not apply 5 days before harvest. Peaches, Apricots, Nectarines. To control Peach Twig Borer, Plum Curculio, June Beetles, San Jose Scale. Do not apply to Apricots and Nectarines within 3 days of harvest, Peaches 1 day. For Peach Tree Borer, apply 2 or 3 sprays to trunk, from ground to scaffold limbs, timed with moth flight.

Cherries, Plums, Prunes to control Mealy Plum Aphids, Plum Curculio, Prune Leaf Hopper, Black Cherry Aphid, San Jose Scale. Do not apply 1 day before harvest.

LAWNS — To control Sod Webworm (Lawn Moth), Earwig, Ant, Flea, Leafhopper, Mosquitoes. Best results, mow lawn, water area well. Do not water for 2 days after application. Repeat 2 weeks later if necessary. Use 4 tablespoons per gal. water per 100 sq. ft.

AROUND HOME, Patio, Outdoor Area to control Ant, Earwig, Flea, Mosquito, Grasshopper. Spray surface. Do not apply to plants. Do not use more than twice a week. Use 4 tbs. per gal. of water.

CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Do not take internally. Skin contact may be harmful. Avoid skin contact, wash hands and face before eating or smoking. Wear clean clothing. Avoid contamination of food. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use on pets or for other uses not covered by this label. Do not leave mixed spray in sprayer. Do not spray indoors. Do not contaminate water by cleaning of equipment or disposal of wastes. Rinse equipment and dispose of waste and empty containers by burying in non-crop lands away from water supplies. Do not reuse containers.

FLAMMABLE — KEEP AWAY FROM HEAT OR OPEN FLAME.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

CHACON CHEMICAL CORPORATION

South Gate, California 90280
A Subsidiary of Leisure
Enterprises, Inc.

EPA REG. NO. 5719-35

CHACON

SEVIN LIQUID GARDEN SPRAY

For Control of Certain Insects on

VEGETABLES - ORNAMENTALS FLOWERS - FRUITS - LAWNS



ACTIVE INGREDIENTS:
Carbaryl (1-naphthyl N-methylcarbamate) 5%

INERT INGREDIENTS 95%

U.S. Patent Nos. 2,903,478 - 3,009,855
*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions.

NET 32 FL. OZ. (1 QUART)

CHACON SEVIN LIQUID GARDEN SPRAY

ORNAMENTALS — Roses, Flowers, Shrubs, Trees, Woody Plants. For the control of Rose Aphids; Japanese, Blister, Elmleaf, Flea, June Beetles; Boxelder, Mealy, Lace, Plant Bugs; Leafhoppers; Leaf Rollers; Psyllid; Exposed Thrips. Gypsy Moth, Scale Insect, Oak Leaf Miners, Tent and Puss Caterpillars, Rose Slug, Orange Tortrix, Cutworms. Use 2 tbs. per gal. of water. Start application when insects or damage appears. Repeat weekly or as needed.

VEGETABLES — Use 2 tbs. per gal. of water. Apply when insects or damage appears. Repeat at 7 to 10-day intervals or as necessary. Tomato, Egg Plant, Peppers to control Tomato Hornworm; Tomato Fruitworm; Lace, Stink and Tarnish Plant Bugs; Leafhoppers. Use up to day of harvest.

Cucumber, Squash, Pumpkin, Melon to control Pickleworm, Melonworm, Cucumber Beetle, Flea Beetles, Leafhoppers, Squash Bugs. Use up to day of harvest.

Lettuce, Carrot, Cabbage, Broccoli, Cauliflower, Spinach, Brussels Sprout, Kohlrabe, Swiss Chard, Radish, Beet, Turnip to control Cabbage Worm and Looper, Armyworms, Flea Beetle, Leafhopper, Cutworm, Corn Earworm. Do not apply within 3 days of harvest. Do not apply within 14 days of harvest in leaf Lettuce, Spinach, and Swiss Chard. Do not treat Beets and Turnips within 14 days of harvest when tops are to be used as food.

Bean, Pea, Corn to control Leafhopper, Grasshopper, Armyworm; Stink and Tarnished Plant Bugs; Cutworm, Japanese Flea and Cucumber Beetles. Use up to day of harvest.

Strawberries, Boysenberries, Raspberries, Blackberries to control Japanese Beetle, Leaf Roller, Leafhopper, Strawberry Weevil. Do not apply within 7 days of harvest on Boysenberries, Raspberries, Blackberries, one day on Strawberries.

FRUITS AND NUTS — Use 2 tbs. per gal. water. Spray when insects or damage appears. Repeat at weekly intervals or as needed. Walnuts to control Codling Moth, Apples, Pears to control Apple Aphid, Pear Slug, Pear Psylla, and Codling Moth. Do not apply to Apples. Pears until 30 days after bloom, or 1 day before harvest on Apples and Pears.

