



1088  
p n 2  
5719-35



ACCEPTED  
SEP 10 1984  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 5719-35

SEVIN\*  
LIQUID GARDEN SPRAY

Aphids      Beetles      Fleas      Hornworms      Ants

VEGETABLES-ORNAMENTALS  
FLOWERS-FRUITS-LAWNS

ACTIVE INGREDIENTS:  
Carbaryl (1-naphthyl N-methylcarbamate) ..... 5.0%  
INERT INGREDIENTS ..... 95.0%  
Total 100.0%

\* Trademark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate.

EPA REG. NO. 5719-35      EPA EST. NO. 5719-CA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION      NET 1 PINT (0.473 l.)

See back panel for additional precautionary statements.

## SEVIN\* LIQUID GARDEN SPRAY

## PRECAUTIONARY STATEMENTS

## Hazards to Humans and Domestic Animals

## CAUTION

Keep out reach of children. Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. Avoid contamination of food or feed.

Statement of practical treatment: If swallowed, do not induce vomiting. Get immediate medical attention. If inhaled, remove affected person from exposure. If in eyes, flush with plenty of fresh water for at least 15 minutes. Get medical attention if irritation persists. If on skin, wash affected area with soap and water.

Note to Physician: Carbaryl is a reversible cholinesterase inhibitor. Atropine is antidotal. Do not administer 2-PAM, opiates or cholinesterase inhibiting drugs.

## Environmental Hazards

This pesticide is toxic to aquatic organisms. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

## Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read the entire label thoroughly. Apply this product only as specified on this label. Do not use inside the home.

For best results use when the air is still, as a thorough cover spray. Apply when insects or damage appears. Repeat at 7-10 day intervals or as necessary.

ORNAMENTALS, VEGETABLES, FRUITS AND NUTS: Use at the rate of 1 fl. oz. (2 tbs.) per gallon of water.

LAWNS, PATIOS, and OUTDOOR AREAS: Use at the rate of 2 fl. oz. (4 tbs.) per gallon of water per 100 sq. ft.

ROSES, FLOWERS, SHRUBS, TREES, WOODY PLANTS: To control Rose Aphids; Japanese, Blister, Elm Leaf, Flea or June Beetles; Boxelder, Mealy, or Lace Plant Bugs; Leafhoppers;

EBB-BB

Leafrollers; Psyllid, Thrips (exposed), Gypsy Moths, Scale Insects, Oak Leafminers, Tent or Puss Caterpillars, Rose Slugs, Orange Tortrix, Cutworms.

Leafminer, Tent or Puss Caterpillar, Rose Slug, Orange Tortrix, Cutworm.

TOMATOES, EGGPLANTS, PEPPERS: To control Tomato Hornworms; Tomato Fruitworms; Lace, Stink or Tarnished Plant Bugs; Leafhoppers. Use up to day of harvest.

CUCUMBERS, SQUASH, PUMPKINS, MELONS: To control Pickleworms; Melonworms; Cucumber or Flea Beetles; Leafhoppers; Squash Bugs. Use up to day of harvest.

LETTUCE, CARROTS, CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS, KOHLRABI, RADISHES, BEETS, TURNIPS: To control Cabbage Worms, Armyworms, Cutworms, Corn Earworms, Loopers, Flea Beetles or Leafhoppers. Do not apply within 3 days of harvest. Do not apply within 14 days of harvest on Leaf Lettuce, Spinach, and Swiss Chard. Do not treat Beets and Turnips with in 14 days of harvest, when tops are used as food.

BEANS, PEAS, CORN: To control Leafhoppers; Grasshoppers; Armyworms; Cutworms; Stink or Tarnished Plant Bugs; Japanese, Flea or Cucumber Beetles. Use up to day of harvest.

STRAWBERRIES, BOYSENBERRIES, RASPBERRIES, BLACKBERRIES: To control Japanese Beetles, Leafrollers, Leafhoppers, Strawberry Weevils. Do not apply within 7 days of harvest on Boysenberries, Raspberries, and Blackberries; 1 day on Strawberries.

PEACHES, APRICOTS, NECTARINES: To control Peach Twig Borers, Plum Curculios, June Beetles, San Jose Scales. Do not apply to Apricots and Nectarines within 3 days of harvest; 1 day on Peaches. For Peach Tree Borers, apply 2 to 3 sprays to trunk, from ground to scaffold limbs, timed with Moth flight.

CHERRIES, PLUMS, AND PRUNES: To control Mealy Plum or Black Cherry Aphids, Plum Curculios, Prune Leafhoppers, San Jose Scales. Do not apply within one day of harvest.

APPLES, PEARS: To control Apple Aphids, Pear Slugs, Pear Psylla, Codling Moths. Do not apply to Apples and Pears until 30 days after bloom, or within 1 day before harvest.

CITRUS: To control Red, Citricola, or Brown Soft Scales. Start spraying in spring. Do not apply within 5 days of harvest.

WALNUTS: To control Codling Moths.

LAWNS: To control Sod Webworms (lawn moths), Earwigs, Ants, Fleas, Leafhoppers, Mosquitoes. For best results, mow lawn, and before application water area well. Do not water for 2 days after application. Repeat 2 weeks later if necessary.

PATIOS & OUTDOOR AREAS: To control Ants, Earwigs, Fleas, Mosquitoes, or Grasshoppers. Spray surface. Do not apply to plants. Do not use more than twice per week.

4. A. 8'

## STORAGE AND DISPOSAL

Storage: Store in original container in a cool place away from sparks or open flame. Keep out of reach of children.  
Disposal: Securely wrap original container in several layers of paper and discard in trash. Do not reuse container.

Notice: Buyer assumes all risk of use, storage and handling of this material not in strict accordance with directions given herewith.

Manufactured By  
CHACON CHEMICAL CORPORATION  
2600 Yates Avenue - City of Commerce, CA 90040  
A GardenAmerica Company.

## LAWN AND GARDEN INSECT CONTROL

### PRECAUTIONARY STATEMENTS

#### Hazards to Humans and Domestic Animals

##### CAUTION

Keep out reach of children. Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. Avoid contamination of food or feed.

Statement of practical treatment: If swallowed, do not induce vomiting. Get immediate medical attention. If inhaled, remove affected person from exposure. If in eyes, flush with plenty of fresh water for at least 15 minutes. Get medical attention if irritation persists. If on skin, wash affected area with soap and water.

Note to Physician: Carbaryl is a reversible cholinesterase inhibitor. Atropine is antidotal. Do not administer 2-PAM, opiates or cholinesterase inhibiting drugs.

#### Environmental Hazards

This pesticide is toxic to aquatic organisms. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

#### Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

#### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read the entire label thoroughly. Apply this product only as specified on this label.

For best results use when the sun is shining and a thorough cover spray. For ORNAMENTALS, VEGETABLES, FRUITS and NUTS, use at the rate of 1 fl. oz. of this per gallon of water. Use at the rate of 2 fl. oz. of this per gallon of water for LAWNS, PATIOS, and OUTDOOR AREAS. Apply when insects or damage appears. Repeat at 7-10 day intervals or as necessary.

ROSES, FLOWERS, SHRUBS, TREES, WOODY PLANTS: To control Rose Aphids; Japanese, Blister, Elmleaf, Flea or June Beetle; Boxelder, Mealy, or Lace Plant Bug; Leafhopper, Leafroller,

Psyllid, Thrips (exposed), Gypsy Moth, Scale Insect, Oak Leafminer, Tent or Puss Caterpillar, Rose Slug, Orange Tortrix, Cutworm.

**TOMATO, EGGPLANT, PEPPER:** To control Tomato Hornworm, Tomato Fruitworm, Lace, Stink or Tarnished Plant Bug; Leafhopper. Use up to day of harvest.

**CUCUMBER, SQUASH, PUMPKIN, MELON.** To control Pickleworm, Melonworm; Cucumber or Flea Beetle; Leafhopper, Squash Bug. Use up to day of harvest.

**LETTUCE, CARROT, CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSELS SPROUTS, KOHLRABI, RADISH, BEET, TURNIP:** To control Cabbage Worm, Armyworm, Cutworm, Corn Earworm, Looper, Flea Beetle or Leafhopper. Do not apply within 3 days of harvest. Do not apply within 14 days of harvest in leaf Lettuce, Spinach, and Swiss Chard. Do not treat Beets and Turnips within 14 days of harvest, when tops are used as food.

**BEANS, PEAS, CORN:** To control Leafhopper, Grasshopper, Armyworm, Cutworm; Stink or Tarnished Plant Bug; Japanese, Flea, or Cucumber Beetle. Use up to day of harvest.

**STRAWBERRIES, BOYSENBERRIES, RASPBERRIES, BLACKBERRIES:** To control Japanese Beetle, Leafroller, Leafhopper, Strawberry Weevil. Do not apply within 7 days of harvest on Boysenberries, Raspberries, and Blackberries; 1 day on Strawberries.

**PEACHES, APRICOTS, NECTARINES:** To control Peach Twin Borer, Plum Curculio, June Beetle, San Jose Scale. Do not apply to Apricots and Nectarines within 3 days of harvest; 1 day on Peaches. For Peach Tree Borer, apply 2 to 3 sprays to trunk, from ground to scaffold limbs, timed with Moth flight.

**CHERRIES, PLUMS, AND PRUNES:** To control Mealy Plum or Black Cherry Aphids, Plum Curculio, Prune Leafhopper, San Jose Scale. Do not apply within one day of harvest.

**APPLES, PEARS:** To control Apple Aphids, Pear Slug, Pear Psylla, Codling Moth. Do not apply to Apples and Pears until 30 days after bloom, or 1 day before harvest.

**CITRUS:** To control Red, Citricola, or Brown Soft Scale. Start spraying in Spring. Do not apply within 3 days of harvest.

**WALNUTS:** To control Codling Moths.

**LAWNS:** To control Sod Webworm (lawn moth), Earwig, Ant, Flea, Leafhopper, Mosquito. For best results, mow lawn, water area well. Do not water for 2 days after application. Repeat 1 week later if necessary.

**PATIOS & OUTDOOR AREAS:** To control Ant, Earwig, Flea, Mosquito, or Grasshopper. Spray surface. Do not apply to plants. Do not use more than twice per week.

#### STORAGE AND DISPOSAL

**Storage:** Store in original container in a cool place away from sparks or open flame. Keep out of reach of children.

**Disposal:** Securely wrap original container in several layers of paper and discard in trash. Do not reuse container.

DEFENDANT COPY

Notice: Buyer assumes all risk of use, storage and handling of this material not in strict accordance with directions given herewith.

Manufactured By  
CHACON CHEMICAL CORPORATION  
2600 Yates Avenue - City of Commerce, CA 90040  
A GardenAmerica Company.