H ... 7

NOV 5 SEP

Gentlumen:

Subject: Chacon's Sevin Garden Dust EPA Degistration Number 5719-36 Your Application Dated September 27, 1982

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. A stamped copy is enclosed for your records.

Sincerely yours,

Jay S. Ellenberger Product Manager (12) Insecticide-Redesticide Branch Registration Division (25-767)

Enclosure

(

TS-767:Ellenberger:DCR-28956:WANG-0945C:jad:Raven:479-2013:11/1/82

UPDATE: DRAFT Sept. 27, 1982



Á

K

5.87

CENT AVAILURE CUPY

6087

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - Harmful if swallowed or inhaled. Avoid breathing of dust. Avoid contact with skin. Avoid contamination of feed and food stuff. Keep out of reach of children.

Wear regular long sleeved clothing during application. Wash exposed skin with soap and water after use.

STATEMENT OF PRACTICAL TREATMENT: If swallowed, promptly drink a large quantity of water and induce vomitting. Get medical attention at once. In case of contact, immediately flush eyes with plenty of fresh water. For skin, wash with plenty of soap and water. Get medical attention if irritation persists.

Note to physicians: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTA' HAZARDS

This pesticide is toxic to aquatic organisms. Do not apply directly to water. Do not apply where run -off is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. apply this product only as specified on this label. For best results apply when air is still. Dust v per and lower leaf surfaces. Apply to soil around the plant. Apply when insects first appear. Repeat as necessary.

Do not apply to Boston ivy or Virginia creeper. Injury to tender foliage may occur if plants are wet when treated or in the presence of high humidity. ORNAMENTALS: Roses, Flowers, Shrubs, Trees, Woody Plants. For Japanese beetles, elm leaf fleas, June beetles, boxelder bugs, mealybugs, lace bugs, tarnished plant bugs, leafhoppers, leaf rollers, thrips (exposed), tent caterpillars, cutworms, earwigs.

VEGETABLES: Tomato, Eggplant, Peppers, To control tomato hornmworm, tomato fruitworm, lace, stink and tarnished plant bugs, leafhopper. Use up to day of harvest.

Cucumber, Squash, Pumpkin, Melon. To control melonworm, cucumber beetle, fleabectle, leafhopper, squash bug. Use up to day of harvest.

Head Lettuce, Carrot, Cabbage, Broccoli, Cauliflower, Brussels Sprouts, Kohlrabi, Radish, Beet, Turnif. To control armyworm, flea beetle, leafhoposm; cutworm, corn earworm. Do not apply within 3 days of harvest (except carrots, up to day of harvest). Do not apply within 14 days of harvest on Leaf lettuce; Spinach and Swiss Chard. Do not treat Beets and Turnips within 14 days of harvest when tops are to be used as food.

Bean, Pea, Corn. To control Leafhopper, grasshopper, armyworm, stink and tarnished plant bugs, cutworm, Japanese, flea and cucumber beetles. Use up () to day of harvest.

Strawberries, Boysenberries, Raspberries, Blackberries. To control Japanese' bectles, leaf roller, leafhopper, strawberry weevil. Do not apply within days of harvest.

FRUITS AND NUTS: Apples, Pears, Walnut. To control apple aphid, pear psying, codling moth. Do not apply to apples until 30 days after bloom or 1 day before harvest on apples and pears.

Peaches, Apricots, Nectarines. To control June beetle, leaf roller, plum curculio, codling moth. Do not apply to apricots and nectarines within 3 days of harvest, peaches 1 day.

Cherries, Plum, Prunes. To control mealy plum aphid, black cherry aphid, plum curculio, prune leafhopper, leaf roller. Do not apply within one day of harvest.

FOR USE ON DOGS AND CATS: To control lice, fleas ticks. Dust liberally over animal and rub thoroughly into skin. For brown dog ticks be sure to also apply on legs, feet and between toes. Comb out dead ticks a few hours after treatment. Keep dust out of eyes.

Do not use on puppies or kittens under four weeks of a ;e.

IN AND AROUND DOG AND CAT QUARTERS: To control lice, leas, ticks. dust liberally in and around sleeping quarters, kennels and dog houses. Force into cracks and crevices. Repeat at weekly intervals or as needed.

STORAGE AND DISPOSAL (12 point)

STORAGE: Do not store product in a high humidity environment. Keep out of reach of children.

DISPOSAL: Do not reuse empty container. Wrap container and put in trash collection.

NOTICE: Buyer assumes all risk of use, storage and handling of this material not in accordance with directions given herewith.

Note: This package is sold by weight and package is full when packed, but due to fluffy nature, the material is inclined to settle.

Manufactured for

CHACON CHEMICAL CORPORATION 2600 Yates Avenue . City of Commerce, California 90040 A Subsidiary of Leisure Enterprises, Inc.

