

10857
17m'2
5719-36



SEVIN[®]
DUST

VEGETABLES
ROSES-ORNAMENTALS
flea control for dogs & cats

EARWIGS CATERPILLAR CUTWORMS
FLEAS
BEETLES
HORNWORMS

ACTIVE INGREDIENT
Carbaryl (1-naphthyl N-methylcarbamate)..... 5%
INERT INGREDIENTS95%

*Trade Mark of Union Carbide Corporation for the active ingredient,
1 naphthyl N-methylcarbamate

ACCEPTED

MAR 5 1984

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 5719-36

EPA Reg. No. 5719-36


EPA Est. No. 5719-CA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

Net Weight 11lb. (0.454 Kg.)

See back panel for additional precautionary statements.



BEST AVAILABLE COPY

SEVIN DUST

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin. Avoid contamination of food or feed.

Statement of Practical Treatment: If swallowed, induce vomiting at once. Get medical attention. If inhaled, remove affected person from exposure. If in eyes, flush with fresh water for at least 3 minutes. Get medical attention if irritation persists. If on skin, wash affected area with soap and water.

Note to Physician Carbaryl is a moderate cholinesterase inhibitor. Atropine is antidotal. Do not administer 2-PAM.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water. Do not apply when runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Physical or Chemical Hazards

Avoid dispersion of dust. Avoid contamination with mineral acids or alkalis.

DIRECTIONS FOR USE


It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read label thoroughly.

Apply this product only as specified on this label. Apply as a light uniform dust spray to the entire plant surface, especially on the underside of foliage. Apply to soil around the plant. Apply when pests first appear. Repeat as necessary. For best results, dust in early morning or late evening when air is still.

ROSES, FLOWERS, SHRUBS, TREES, WOODY PLANTS: To control Japanese or June Beetle; Boxelder, Lace or Tarnished Plant Bug; Mealybugs, Elm Leaf Flea, Leafhoppers, Leaf Rollers, Thrips (exposed), Tent Caterpillar, Cutworms, Earwigs.

TOMATO, EGGPLANT, PEPPER: To control Tomato Fruitworm or Hornworm; Lace Stink or Tarnished Plant Bug; Leafhoppers. Use up to day of harvest.

CUCUMBER SQUASH, PUMPKIN, MELON: To control Melonworm, Cucumber or Flea Beetle; Leafhoppers, Squash Bug. Use up to day of harvest.



LETTUCE, CARROT, CABBAGE, BROCCOLI, CAULIFLOWER, BRUSSEL SPROUTS, KOHLRABI, RADISH, BEET, TURNIP: To control Armyworms, Cutworms, Corn Earworm, Flea Beetle, Leafhoppers. Do not apply within 3 days of harvest; 1-day on Carrots. Do not apply within 14 days of harvest on Leaf Lettuce, Spinach and Swiss Chard. Do not treat Beets and Turnips within 14 days of harvest when tops are to be used as food.

BEANS, PEAS, CORN: To control Leafhoppers, Grasshoppers, Armyworms, Cutworms; Stink or Tarnished Plant Bug; Japanese, Flea or Cucumber Beetle. Use up to day of harvest.

STRAWBERRIES, BOYSENBERRIES, RASPBERRIES, BLACKBERRIES: To control Japanese Beetle, Leaf Rollers, Leafhoppers, Strawberry Weevil. Do not apply within 7 days of harvest; except Strawberries, 1-day.

PEACHES, APRICOTS, NECTARINES: To control June Beetle, Leaf Roller, Plum Curculio, Codling Moth. Do not apply to Apricots and Nectarines within 3 days of harvest; 1-day on Peaches.

CHERRIES, PLUMS, PRUNES: To control Mealy Plum or Black Cherry Aphid; Plum Curculio, Prune Leafhopper, Leaf Roller. Do not apply within 1-day of harvest.

APPLES, PEARS, WALNUTS: To control Apple Aphid, Pear Psylla, Codling Moth. Do not apply to Apples and Pears until 30 days after bloom, or 1-day before harvest.

FOR USE ON DOGS AND CATS: To control Lice, Fleas, Ticks. Dust liberally over animal and rub thoroughly into skin. For Brown Dog Ticks be sure to also apply on legs, feet and between toes. Comb out dead ticks a few hours after treatment. Keep dust out of eyes. Do not use on puppies or kittens under four weeks of age.

IN AND AROUND DOG AND CAT QUARTERS: To control Lice, Fleas, Ticks. Dust liberally in and around sleeping quarters, kennels and dog houses. Force into cracks and crevices. Repeat at weekly intervals or as needed.

STORAGE AND DISPOSAL

Storage: Store in original container in a dry place. Do not contaminate food or feed. Keep out of reach of children.

Disposal: Securely wrap original container in several layers of paper and discard in trash. Do not re-use container.

Notice: Buyer assumes all risk of use, storage and handling of this material not in strict accordance with directions given herewith.

Manufactured For
 CHACON CHEMICAL CORPORATION
 2600 Yates Avenue - City of Commerce, CA 90540
 A GardenAmerica Company

BEST AVAILABLE COPY