


CHACON

DIAZINON GARDEN DUST

For Control of Certain Named Insects on:

VEGETABLES-FRUITS-FLOWERS
ORNAMENTALS-LAWNS

Preplant Soil Insecticide for Vegetables



Earwig

Flea

Cutworm

Beetle

ACTIVE INGREDIENTS:

0,0-diethyl 0,0-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate	40
INERT INGREDIENTS	96.0
TOTAL	100.0

DIAZINON is a Registered Trademark of
CIBA-GEIGY CORP.

EPA REG. NO. 5719-37

CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions.

NET WEIGHT: 16 OZ. (1 LB.)

CHACON DIAZINON GARDEN DUST

Read the entire label carefully. Use only as directed in accordance with label conditions and directions. For best results apply in early morning or evening when air is still.

DIRECTIONS

VEGETABLES AND FRUITS. Dust lightly on first appearance of insect or damage. Make a complete coverage of all pests and foliage. Repeat applications as necessary for control, but observe the number of days before harvest for last application noted by each crop (for example, on Beans, you may apply this dust up to 7 days before harvest.)

Beans, for the control of Black Bean Aphids, Mexican Bean Beetles, Leafminers, Cucumber Beetles. Do not apply within 7 days of bean harvest. Broccoli, Beets, Cabbage and Cauliflower, Radishes, for control of Aphids, Diamondback Moths, Imported Cabbageworms. Make applications at 7-10 day intervals. Do not apply to Broccoli or Cauliflower within 5 days of harvest or to Cabbage within 7 days of harvest. Do not apply to Beets within 14 days of harvest. Do not apply to Radishes within 10 days of harvest.

Grapes for control of Pacific Spider Mites, Leafhoppers, Grape Berry Moths, Grape Leaf Folders. Repeat as necessary. Do not apply within 18 days of harvest. Lima Beans for control of Black Bean Aphids, Mexican Bean Beetles, Dipterous Leafminers, Leafhoppers, and Mites. Do not apply within 7 days of harvest. Squash and Melons for the control of Aphids, Leafhoppers, Mites, Serpentine Leafminers, and Thrips. Repeat as necessary. Do not apply to melons and Winter Squash within 3 days of harvest, or to Summer Squash within 7 days of harvest. Onions, for control of Onion Thrips. Do not apply within 10 days of harvest. Peas, for control of Aphids and Leafminers. Repeat as necessary. Peas may be harvested, immediately. Peppers for control of Aphids, Serpentine Leafminers. Do not apply within 5 days of harvest. Potatoes for the control of Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous Leafminers, Leafminers. Do not apply within 14 days of harvest. Strawberry, for the control of Aphids, Two-Spotted Mites, Cyclamen Mites, Leafrollers. Do not apply within 5 days of harvest. Sweet Corn, for the control of Corn Earworms, apply when silks first appear. Two to three repeat applications at 2-3 day intervals will be necessary. Do not apply within 2 days of harvest. Grasshoppers, apply when insects appear. Repeat as necessary. Tomatoes for the control of Aphids, Dipterous Leafminers, Vinegar Flies (*Phytophila* spp.), Banded Cucumber Beetles, Fall Armyworms, Southern Armyworms. Repeat as necessary. For Leafminers and vinegar flies, apply at weekly intervals or as required. Do not apply to tomatoes within 3 days of harvest. Apples and Pears for the control of Aphids and Mites. Do not apply within 14 days of harvest.

FLOWERS AND ORNAMENTALS: Azalea, Camellia, Carnation, Chrysanthemum, Dahlias, Gladioli, Rhododendron, Rose and Arbutus, Boxwood, Elm, Juniper, Oak, Pine, Spruce for the control of Aphids, Mites, Leafhoppers, Mealy Bugs, White Flies, Scale Crawlers, Lace Bugs. Do not use on certain ferns, poinsettias, hibiscus, papaya, pilea, and gardenia, since injury may occur. Not effective against certain spider mites on evergreens. Do not apply when honey bees are foraging in the garden.

LAWNS: Bluegrass, Merion, Fescue, St. Augustine, Dichondra, Bent and Bermuda for the control of Ants, Armyworms, Chinch bugs, cutworms, Earwigs, Fleas, Sod Webworms. Apply this package (1 lb.) evenly as a broadcast over 250 square feet. Water lawn thoroughly after application. Repeat application in 7-10 days if necessary.

PREPLANT SOIL INSECTICIDE prior to planting the following vegetables: Bean, Broccoli, Cabbage, Cauliflower, Corn, Cucumbers, Endive, Lettuce, Melons, Onions, Parsnips, Peas, Peppers, Potatoes, Spinach, Squash and Tomatoes. To control Cutworms, Seed corn maggots, and wireworms. Apply evenly as a broadcast treatment over the surface of the soil. The amount in this package (1 lb.) should be broadcasted over 150 square feet. Immediately after spreading, mix thoroughly with the soil to a depth of 6-8 inches.

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Do not take internally. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after handling. Do not store near or contaminate feed or foodstuffs. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label. Do not reuse container. Destroy when empty.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given here-with.

EPA EST. 5719-CA 1

CHACON CHEMICAL CORPORATION
5245 Chakemco Street • South Gate, California 90280
A Subsidiary of Leisure Enterprises, Inc.