

E
CHACON

DIAZINON
DUST


VEGETABLES
LAWNS - GARDENS

ALSO A PREPLANT SOIL
INSECTICIDE FOR VEGETABLES



ANTS

CUTWORMS



EARWIGS



APHIDS



BEETLES

ACTIVE INGREDIENT

Diethyl 0-0-2 isopropyl 0 methylid
pyrimidyl phosphorothioate
INERT INGREDIENTS
TOTAL

40%
96.0%

© 1974, Chacon & Register, Fresno, Calif.
TRADE MARK

ACCEPTED
with conditions
by EPA on 3/1/84

MAR 3 1984

Under the Insecticide,
Fungicide, and Rodenticide Act
as amended, this pesticide
registered under Reg. No.

PRECAUTIONARY STATEMENT

PRECAUTIONARY STATEMENT

Hazards to Humans and Domestic Animals

Keep out of reach of children. Harmful if swallowed. Avoid inhalation of dust. Avoid contact with skin. Avoid contamination of food or feed.

Statement of Practical Treatment: If swallowed, induce vomiting at once. Get medical attention. If inhaled, remove affected person from exposure. If in eyes, flush with fresh water for at least 3 minutes. Get medical attention if irritation persists. If on skin, wash affected area with soap and water.

Environmental Hazards

This product is toxic to fish. Do not apply directly to water. Do not apply when runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Physical or Chemical Hazards

Avoid dispersion of dust. Avoid contamination with mineral acids or alkalis.

DIRECTION FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label thoroughly. Apply this product only as specified on this label.

Apply as a light uniform dust spray to the entire plant surface, especially on the underside of foliage. Apply to soil around the plant. Apply when pests first appear. Repeat as necessary. For best results, apply in early morning or late evening.

BEANS: To control Aphis, Mexican Bean or Cucumber beetle; Leafminer. Do not apply within 7 days of harvest.

BROCCOLI, BRASSICA VEGETABLES, CAULIFLOWER, RADISHES: To control Aphids, Imported Cabbageworm. Apply at 7-10 day intervals. Do not apply within the following number of days before harvest: 5 days-Broccoli, Cauliflower; 7 days-Cabbage; 10 days-Radishes; 14 days-Beets.

LIMA BEANS: To control Aphis, Bean Aphid, Mexican Bean beetle, Dipterous Leafminer, Leafhoppers, Mites. Do not apply within 7 days of harvest.

SQUASH, MELONS: To control Aphids, Leafhoppers, Mites, Serpentine Leafminer, Thrips. Do not apply within the following number of days before harvest: 3 days-Melons, Winter Squash; 7 days-Summer Squash.

BEST DOCUMENT AVAILABLE

ONIONS: To control Onion Thrips. Do not apply within 10 days of harvest.

LEAS: To control Aphids, Leafminers. Use up to day of harvest.

FEIPEK: To control Aphids, Serpentine Leafminer. Do not apply within 7 days of harvest.

POTATOES: To control Aphids; Flea or Colorado Potato Beetle; Dipperous Leafminer, Leafminers. Do not apply within 14 days of harvest.

SWEET CORN: To control Corn Earworm, apply when silks first appear. Two to three repeat applications will be necessary at 2-3 day intervals.

To Control Grasshoppers, apply as necessary. Do not apply within 2 days of harvest.

TOMATOES: To control Aphids, Dipperous Leafminer, Vinegar Fly (*Drosophila* spp.), Banded Cucumber Beetle; Fall or Southern Armyworm. Do not apply within 3 days of harvest.

APPLES, PEARS: To control Aphids, Mites. Do not apply within 14 days of harvest.

GRAPES: To control Pacific Spider Mite, Leafhoppers, Grape Berry Moth or Leaf Folder. Do not apply within 18 days of harvest.

AZALEAS, CAMELLIAS, CHRYSANTHEMUMS, DAHLIAS, GLADIOLI, RHODODENDRONS, ROSES, ARBORVITAE, BOXWOOD, ELM, JUNIPER, OAK, PINE,

SPRUCE: To control Aphids, Mites, Leafhoppers; Mealy or Lace Bug;

White Fly, Scale Crawler, Do not use on certain Ferns, Poinsettias, Hibiscus, Papaya, Pilea, Gardenia, since injury may occur. Not effective against certain Spider Mites on evergreens.

BLUEGRASS, MERION, FESCUE, ST. AUGUSTINE, DICHONDRA, BENT, AND

BERMUDA LAWNS: To control Ants, Armyworms, Cutworms, Sod Webworm, Chinch Bug, Earwigs, Fleas. Apply at the rate of 1 lb. evenly as a broadcast over 250 sq. ft. (10 oz. per 155 sq. ft.).

Water thoroughly after application. Repeat as necessary at 7-10 day intervals.

AS A PREPLANT SOIL INSECTICIDE IN BEANS, BROCCOLI, CABBAGE, CAULIFLOWER, CORN, CUCUMBERS, ENDIVE, LETTUCE, MELONS, ONIONS, PARSNIPS, PEAS, PEPPERS, POTATOES, SPINACH, SQUASH, TOMATOES.

To control Cutworms, Seed-corn Maggot, Wireworms. Apply evenly as a broadcast soil treatment at the rate of 1 lb. per 450 sq. ft. of area (10 oz. per 250 sq. ft.). Mix thoroughly with the soil to a depth of 6 to 8 inches after application.

STORAGE AND DISPOSAL

Storage: Store in original container in a dry place. Do not contaminate food or feed. Keep out of reach of children.

Disposal: Completely wrap or seal in plastic. Place in layers of paper and discard in trash. Do not reuse containers.

Precautions: Avoid contact with eyes, storage and handling of this material should be strictly in accordance with directions on label and MSDS.

Manufactured by
CHACON CHEMICAL CORPORATION
2600 Yates Avenue - City of Commerce, CA 90040
A GardenAmerica Company

BEST DOCUMENT AVAILABLE

300/116172
17/3

6/10

Mr. Reynaldo Sanabrais
Chacon Chemical Corporation
2600 Yates Avenue
City of Commerce, CA 90040

Dear Mr. Sanabrais.

Subject: Amendment - Update Label
Chacon Diazinon Dust
EPA Registration No. 5719-37
Your Application Dated February 13, 1984

The amendment referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) is acceptable, provided that you make the labeling changes listed below before you release the product for shipment bearing the amended labeling:

1. Add "Diazinon" in front of the diazinon formula name which should then appear in brackets.
2. Please refer to the enclosed type size guide for assistance in selecting the correct print size for the front panel signal word, child hazard warning, and the heading "Storage and Disposal."
3. Revise the If Swallowed advice to read:

If swallowed - Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with a finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

4. Revise the first sentence of the Environmental Hazards to read "This product is toxic to fish, birds and other wildlife." Add a second sentence: "Birds feeding on treated areas may be killed." You may delete "Do not apply when runoff is likely to occur" since this is not appropriate for a dust product.

CONCURRENCES							
SYMBOL ▶							
SURNAME ▶							
DATE ▶							

5. Revise the bee precaution to read:

This pesticide is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

Submit five (5) copies of your revised final printed labeling before you release the product for shipment with the amended labeling.

A stamped copy of the label is enclosed for your records.

Sincerely yours,

George T. LaRocca
Product Manager (15)
Insecticide-Rodenticide Branch
Registration Division (TS-767)

Enclosure

RD/DIS:DCR:05466:LaRocca:pjh:Raven:557-2226:CBI-21:2/27/84:Del.3/12/84