

7-19-40

# CHACON

## DU-O-CIDE MULTI-PURPOSE GARDEN SPRAY



EPA REG. NO. 5719-40 AA

EPA EST. No. 5719-CA-1

**FRUIT TREES** — Apples, Pears. For control of Aphids, Thrips (exposed) Blister Mites, Powdery Mildew. Apply as insects or diseases appear. Begin when buds swell. Repeat when fruit buds show color and again at petal fall. Wash fruit thoroughly before eating. Do not spray during blooming period. As a **DORMANT SPRAY** — For control of Leaf Blight, Peach Leaf Curl, Powdery Mildew, Shot Hole Fungus on Nectarines, Peaches, Plums. Use 4 tablespoons (2 oz.) per gallon of water. Apply when leaves begin to fall, again when buds begin to swell in the spring and when buds begin to show color. Be sure to spray all parts of the tree. Keep material in original container. Do not put into food or drink containers. Do not leave mixed spray in sprayer. Do not store diluted spray, make new mixture each time. Do not store unused diluted spray.

**WARNING:** Poisonous if swallowed or absorbed through the skin. Do not breathe spray mist. Do not get in eyes, on skin, or on clothing. In case of contact, wash immediately with soap and water. Wash all contaminated clothing before reusing. Avoid contamination of feed or feedstuffs.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.** This product is toxic to fish, shrimp, and crabs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not reuse container. Destroy it by perforating or crushing. Bury or discard in a safe place away from water supplies. Apply this product only as specified on this label.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**ACTIVE INGREDIENTS:**

Copper Oleate*	15.00%
Pyrethrins	0.40%
Rotenone	1.25%
Other Cube Extractives	1.25%
Light Medium Petroleum Oil**	15.00%
<b>INERT INGREDIENTS</b>	<b>67.10%</b>
<b>TOTAL</b>	<b>100.00%</b>

\*Copper as element, 1.30%  
\*\*Castor Oil, Deodorized, Agricultural Grade, with Petroleum Oil  
Light Medium Unsaponified Residue 42%

**CHACON CHEMICAL CORPORATION**  
2600 Yates Ave. • City of Commerce, CA. 90040  
A Subsidiary of Leisure Enterprises, Inc.

### CHACON DU-O-CIDE MULTI-PURPOSE GARDEN SPRAY

Read entire label carefully before using. Use only in accordance with label warnings and directions.

**DIRECTIONS**

For control of all insects and diseases listed, mix 1 tablespoonful (1/2 oz.) per gallon of water. (Except when using as a **DORMANT SPRAY**, use 4 tablespoons (2 oz.) per gallon.) Note: Do not mix sticking or wetting agents or any other materials with this product. Spray plants thoroughly, including both sides of leaves and around soil at base of plant. May be used on fruits and vegetables listed one day before harvest.

**FLOWERS, SHRUBS, ORNAMENTAL TREES** — (Azaleas, Begonias, Camellias, Chrysanthemums, Grape Myrtle, Euonymus, Dahlias, Delphinium, Fuchsias, Roses, Saururus, Stocks, Sweet Peas. For the control of Aphids, Diabrotica Beetles, Leafhoppers, Mealybugs, Whitefly Larvae, Spider Mites, Red Spider Mite, Hops Slug, Thrips (exposed) Black Spots, Leaf Spots, Powdery Mildew, and Rust. Begin when foliage or diseases appear. Repeat at 7 to 10-day intervals or as necessary.

**VEGETABLES** — Beans, Broccoli, Cabbage, Cauliflower, Kale, Turnips, Cucumbers, Lettuce, Melons, Peas, Tomatoes. For control of Aphids, Diabrotica Beetles, Thrips (exposed), Squash Bugs (nymphs), Leafhoppers, Mealybugs, Powdery Mildew and Rust only on Beans. Apply as diseases or insects appear. Repeat weekly or as needed. Wash vegetables thoroughly before eating. Do not spray during bloom period.

**CITRUS** — Oranges, Lemons, Grapefruit. For control of Aphids, Leafhoppers, Whitefly Larvae, Red Spider Mites, Scale Crawlers, Sooty Mold. Apply as diseases or insects appear, repeat as needed. Do not spray during bloom period.

**WALNUTS** — For control of Aphids, Red Spider Mites, Leaf Spot. Apply after blooms drop. Repeat at two-week intervals or as needed. Do not use after husks begin to crack.

**BERRIES** — Blackberry, Raspberry, Strawberry. For control of Aphids, Red Spider Mites, Leafrollers, Leaf Spot. Apply as insects or disease appear. Repeat as needed. Do not use after fruit begins to form or during bloom. Wash fruit thoroughly before eating.

**WARNING:** KEEP OUT OF REACH OF CHILDREN  
See back panel for additional cautions.

NET 32 FL. OZ. (1 QUART)

ACCEPTED  
5719-40