

Columbia

of Baltimore

AND DECORATED
DOORS, CLOSURES,
AND SPECIALTIES.



Proof

COLUMBIA
SPECIALTY COMPANY, INC.
6301 EASTERN AVE
BALTIMORE, MD. 21224

tainers: 6 tablespoonfuls per gal. of water. Spray around foundation, under porches, along fences, shrubbery and other infested areas.
HOUSEHOLD PESTS, Roaches, Ants, Spiders, Mosquitoes, Silverfish, Crickets, Cloverleafs, Earwigs, in and around homes: 8 tablespoonfuls per 1 gallon of water. Apply coarse spray or brush under sinks, around windows, baseboards, drainboards and other areas. Avoid contamination of food, dishes, utensils and water.

CAUTIONS. Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin, wash skin and hands thoroughly after using. Avoid contamination of feed or food stuffs. Malathion is a cholinesterase inhibitor and can cause symptoms similar to those caused by other organic phosphate compounds. If poisoning should occur, call a physician immediately. Atropine is antidotal.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME. Keep children and pets off treated areas until material is washed into the soil and grass is dry. Do not use on Boston Maiden Hair or Pteris Ferns.

Do not leave mixed spray in sprayer. Do not store diluted spray. Make new mixture each time. Do not contaminate water by cleaning of equipment or disposal of waste. Rinse equipment and dispose of waste and empty containers by burying in non-crop lands away from water supplies. Do not reuse containers. This product will kill fish and wild life. Do not contaminate lakes, ponds, streams, woodlands and other non-crop areas.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

EPA REG. NO 5719-41 AA EPA EST. 5719-CA-1

CHACON CHEMICAL CORPORATION

2600 Yates Avenue • City of Commerce, California 90040
 A Subsidiary of Leisure Enterprises, Inc.

CHACON

**50%
MALATHION**

Active Ingredients:
 Malathion* 50.00%
 Aromatic Petroleum Derivative Solvent 40.37%
 Inert Ingredients 9.63%
 *O,O dimethyl dithiophosphate of diethyl mercaptosuccinate

NET 32 FL. OZ. (1 QUART)

CAUTION: Keep out of Reach of Children - Other Cautions on Back Panel

Read entire label with label cover both necessary. less other agents or
ORNAMENTALS
 santhemum
 spoonfuls
 Mealybugs
 Tarnished
 1 tablespoo
 Oyster Sh
FRUITS, A
 gallon
 (applied)
 Tarnished
 Crawlers: 1
 to Pears w
 Malathion
 Grapes.
CITRUS, G
 White Flie
 posed); 2
 7 days of
VEGETABLES
 Kale, Beet
 water. Ret
 Bean Lea
 tablespoo
 Peas with
 Cauliflow
 bloom per
HOUSE FL

BEST DOCUMENT AVAILABLE

ACCEPTED
 AUG 15 1975
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 this pesticide
 is registered under
 EPA Reg. No. 5719-41

51

ter. Spray around foundation,
y and other infested areas.
nders, Mosquitoes, Silverfish,
around homes: 8 tablespoon-
-arse spray or brush under
rainboards and other areas.
utensils and water.

aid breathing of spray mist.
and hands thoroughly after
r food stuffs. Malathion is a
e symptoms similar to those
mpounds. If poisoning should
ropine is antidotal.

NEAR HEAT OR OPEN FLAME.
eas until material is washed
use on Boston Maiden Hair

Do not store diluted spray.
not contaminate water by
waste. Rinse equipment and
eas by burying in non-crop
not reuse containers.

e. Do not contaminate lakes,
non-crop areas.
exposed to direct treatment
ormation may be obtained
ension Service.


use, storage or handling of
with directions given herewith.

PA EST. 5719-CA-1

L CORPORATION

ommerce, California 90040
e Enterprises, Inc.

CHACON



NET 32 FL. OZ. (1 QUART)

CAUTION: Keep out of Reach of Children - Other Cautions on Back Panel

CHACON 50% MALATHION

Read entire label carefully. Use only as directed in accordance with label cautions, warnings and directions. Spray thoroughly, cover both sides of leaves and other parts of the plant. Repeat as necessary. Can be used up to 3 days of harvest on food crops unless otherwise specified. NOTE: Do not mix sticking or wetting agents or any other spray materials with this product.

ORNAMENTALS, Roses, Azaleas, Gardenias, Camellias, Stocks, Chrysanthemums, and other flowers and shrubs. Aphids: 1½ to 2 teaspoonfuls per gallon of water. Red Spider Mites, White Flies, Mealybugs, Woolly Aphids, Rose Leafhoppers, Thrips (exposed), Tarnished Plant Bug, Box Elder Bug, Japanese Beetles (Adult): 1 tablespoonful per gallon of water. Black Scale, Soft Brown Scale, Oyster Shell Scale: 1½ to 2 tablespoonfuls per gallon of water.


FRUITS, Apples, Pears, Grapes, Aphids: 1½ to 2 teaspoonfuls per gallon of water. Red Spider Mites, Mealybugs, White Flies, Thrips (exposed), Japanese Beetle (Adult), Pear Psyllid, Grape Leafhopper, Tarnished Plant Bug: 1 tablespoonful per gallon of water. Scale Crawlers: 1½ to 2 tablespoonfuls per gallon of water. Do not apply to Pears within 1 day of harvest. Do not spray during bloom period. Malathion may injure the foliage on some varieties of Apples and Grapes.

CITRUS, Orange, Lemon, Lime, Grapefruit, Tangerines, Aphids, White Flies, Black Scale, Purple Scale, Yellow Scale, Thrips (exposed): 2 teaspoonfuls per gallon of water. Do not apply within 7 days of harvest. Do not spray during bloom period.

VEGETABLES, Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Kale, Beans, Peas, Aphids: 1½ to 2 teaspoonfuls per gallon of water. Red Spider Mites, Mealybugs, White Flies, Thrips (exposed), Bean Leafhopper, Japanese Beetle (Adult), Tarnish Plant Bug: 1 tablespoonful per gallon of water. Do not apply to Broccoli and Peas within 3 days of harvest and to Brussels Sprouts, Cabbage, Cauliflower or Kale within 7 days of harvest. Do not spray during bloom period. Do not treat beans within one day of harvest.

HOUSE FLIES (outside only) around Dwellings, Patios, Garbage Con-

BEST DOCUMENT AVAILABLE



51

BEST DOCUMENT AVAILABLE

52