

hibitor. Atropine is antidotal.

WARNING — FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. This product is highly toxic to bees exposed to direct treatment or residuals on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use on pets or other uses not covered by label. Do not leave mixed spray in sprayer. Do not spray indoors. Do not contaminate water by cleaning of equipment or disposal of wastes. Rinse equipment and dispose of waste and empty container by burying in non-crop lands away from water supplies. Do not reuse container.

ACTIVE INGREDIENTS

*Carbaryl (1-naphthyl N-methylcarbamate)	5.00%
**S-(2-(Ethylsulfanyl) ethyl) O, O Dimethyl phosphorothioate	5.00%
***2, 4-Dinitro-6-octyl phenyl crotonate	1.46%
***2, 6-Dinitro-4-octyl phenyl crotonate	
***Nitrooctyl phenols (principally diortho)	0.10%
Aromatic Petroleum Derivative Solvent	29.37%
INERT INGREDIENTS	59.07%

*U.S. Patent Nos. 2,903,478 3,009,895

**U.S. Patent No. 2,791,599

***A mixture of 1-methylheptyl, 1-octylheptyl and 1-propylpentyl isomers.

EPA REG. NO. 5719-45

CHACON FERTILIZER CORPORATION

5245 Chalmers Street • South Gate, California 90280

A Subsidiary of Leisure Enterprises, Inc.



CHACON
SYSTEMIC SPRAY
FUNGICIDE

MULTIPURPOSE INSECTICIDE
with Powdery Mildew Control

FOR ORNAMENTALS
ROSES FLOWERS



CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions.

NET 16 FL. OZ. (1 PINT)

CHACON SYSTEMIC SPRAY WITH FUNGICIDE

Read entire label. Use only in accordance with label cautions, warnings and directions. **DO NOT USE ON FRUITS OR VEGETABLES.**

Water thoroughly all areas to be treated 12 to 24 hours before. Do not spray when foliage of plant is wet.

DIRECTIONS
ORNAMENTALS, ROSES and FLOWERS. Use 2 table-spoonfuls per gallon of water for the control of the following insects. Spray entire plant, applying to upper and underside of leaves.

<u>Aphids</u>	<u>Japanese Beetles</u>	<u>Red Spider Mites</u>
<u>Cutworms</u>	<u>June Beetles</u>	<u>Tent Caterpillars</u>
<u>Earwigs</u>	<u>Lace Bugs</u>	<u>Thrips (Exposed)</u>
<u>Elmleaf Beetles</u>	<u>Leafhoppers</u>	<u>White Flies</u>
<u>Flea Beetles</u>	<u>Leaf Miners</u>	

For the Control of **POWDERY MILDEW** on Roses, Dahlias, Delphiniums, Grape Myrtle. Use 4 table-spoonfuls per gallon water. Protect new growth by spraying at 2-week intervals and apply to upper and underside of leaves.

CAUTION: Harmful if swallowed, inhaled, or absorbed through the skin. Do not take internally. Avoid skin contact. Wash hands and face before eating or smoking. Wear clean clothing. Avoid contamination of food. If swallowed, call physician. **NOTE TO PHYSICIANS:** This product is an organic phosphate which is a cholinesterase inhibitor. Atropine sulphate is antidotal. Carbaryl is a moderate, reversible, cholinesterase in-

ACCEPTED

JUL 5 1973

UNDER THE FEDERAL INSECTICIDE
AND FUNGICIDE ACT
AS AMENDED
REGISTERED UNDER NO. 5719-45 SUBJECT
TO ATTACHED COMMENTS.

5719-45