
in and around GARDEN - LAWN VEGETABLES LISTED





CUTWORM

SOWBUG

TREATS UP TO 1250 SQ. FT.

Active Ingredient
Carbaryl (1 naphthyl N-methylcarbamate)
Inert Ingredients

95%

Total

KEEP OUT OF REACH OF CHILDREN

CAUTION

See additional precautionary statements at end of label

NET WEIGHT 20 OZ. (11/4 LB.)

CHACOH CUTWORM & JUWBUG CONTROL

PM 12-

Directions for Applying Cutworm & Sowbug Control

It is a violation of federal law to use this product in a manner inconsistant with its labeling Chacon Cutworm & Sowbug Control is formulated to attract and kill cutworms, sowbugs, earwigs, crickets, dar' ing ground beetles, grasshoppers, millipedes, armyworms and ants coming in contact with pellets. For use in and around ornamentals, ground covers, lawns, and certain lists vegetables.

ORNAMENTALS, GROUND COYERS, LAWNS: Start treatment at first sign of pest activity. Water area first, then do not water for at least 24 hours. Spread pellets lightly over area Apply at the rate of 1½ pounds per 1250 sq. ft. Sowbugs and Cutworms feed mainly at night. It is best to put out the pellets in late afternoon or evening. Application should be repeated at 2-week intervals as long as needed.

VEGETABLES LISTED: Apply on the ground and adjacent to plants. Do not apply to EDIBLE PARTS OF THE PLANTS, Broadcast applications may be used when there is a heavy infestation. Repeat applications as necessary, Control will be obtained with $\frac{3}{2}$ lbs to $\frac{11}{4}$ lbs per 1250 sq. (L. of bait. For the control of sowbugs, cutworms, armyworms, earwigs, millipedes and ants.

Сгор	Dosage per 1250 Sq. Ft	Days Between Final Application and Harvest
Asparaç is, Strawberries	I 1/4 lbs	1 Day
Beans, Carrots, Corn, Sweet Corn, Eggplant, Okra, Peppers, Potato, Tomato	11/4 lbs	NONE
Cucumbers, Melons, Squash, Peas	₹⁄5 lbs	NONE
Bruccoli, Brussels sprouts, cabbage, cauliflower, head lettuce, garden beet roots, parsnip, radish, rutabaga, turnip (roots)	11/4 lbs	3 Days
Collards, Endive, Garden beet tops, Kale, leaf tettuce. parsley, spinach, Swiss chart turnip tops	1 1/4 1bs	14 Days
		1

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Harmful if inhaled or swallowed Avoid breathing of dust. Do not take internally Avoid contact with skin and eyes. Wear regular long-sleeved work clothing. Change to clean clothing daily Wash hands and face before eating Wash thoroughly after handling. For protection of honeybees avoid unnecessary use during period they are visiting crop.

NOTE for Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or Cholinesterase inhibiting drugs. Avoid contamination of food, feed, water supplies, streams and ponds. Keep pets off treated areas.

ENVIRONMENTAL HAZARDS

This product is toxic to wildlife. Use with care when applying in areas frequented by birds. Birds and other wildlife feeding on treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

STORAGE & DISPOSAL

STORAGE: Store in a cool, dry place

DISPOSAL: Do not re use empty container. Wrap container and put in trash collection.

NOTICE: Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith

EPA Reg. No. 5719-46

EPA Est 2935-CA-4

MANUFACTURED FOR
CHÁCON CHEMICAL CORPORATION
2600 Yates Avenue, City of Commerce, California 90040
A Subsidiary of Leisure Enterprises, Inc.

5714-46

Grader the First cold for the Condens of Breat Library Cold for the Co

NI BEE