

Do not spray within 7 days of harvest. Do not spray on fruit or vegetables from vine that is infected with Tomato Hornworm.

Use 6 tablespoons per gallon of water to control Codling Moth on Apples, Pears and Walnuts. For control of Apple Moth, Apple Maggot, Orange Fruit Moth, or Plum curculio. Apply at petal fall and every 7 to 14 days thereafter until control is achieved. On Rose Beetles, Leaf Rollers, Weevils, on Strawberry, Boysenberry, Raspberry, Blackberry. Do not use within 7 days of harvest except on Strawberry within 3 days.

CAUTION: Highly toxic if swallowed. Absorbed through the skin. Do not take internally. Skin contact may be harmful. Avoid skin contact with eyes, mouth and face before eating or smoking. Wear protective gear to prevent contamination of food.

This product is highly toxic to bees exposed to direct treatment or residues on plants. Protective information may be obtained from your Cooperative Extension Service. Do not use on flowers for which bees are known to visit by this insect. Do not allow spray to drift onto blossoms. Do not spray windbreaks.

This product is toxic to fish. Keep out of lakes, streams, ponds, etc., especially where runoff is likely to occur. Do not spray when weather conditions favor drift from areas treated. Do not spray on areas where spraying of equipment or disposal of wastes. Apply the product at rates specified on this label. Do not reuse containers.

WARNING - FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME
Malathion can be a respiratory irritant and can cause symptoms similar to those caused by organic phosphate compounds. If symptoms develop, seek medical attention immediately. Atropine is an antidote.

NOTE FOR PHYSICIAN Anticholinesterase irreversible inhibitor. Use atropine to reverse the Atropine-like effects.


NOTICE: Please consult the Material Safety Data Sheet of the manufacturer for complete health and safety information.

EPA REG. No. 5719-56-AA EPA EST. No. 5719-CA-1

CHACON CHEMICAL CORPORATION
2600 Yater Ave. • City of Commerce, CA 90040
S. California • U.S.A.

RUGBY

1978



CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions.

NET 32 FL. OZ. (1 QUART)

CHACON FRUIT AND VEGETABLE SPRAY

Read entire label carefully before using. Do not mix with other directions.

DIRECTIONS

Apply when insects or damage occurs. Repeat at 7 to 10 day intervals, as necessary. Do not use more than 10% volume of diluted spray per acre.

VEGETABLES: Use 3 tablespoons per gallon of water to control Leafhopper on Tomatoes and leafy vegetables; on the Worm and Melonworm on Cucumber, Squash, Eggplant, Melon, Cantaloupe on leafy vegetables. Do not use within 7 days of harvest except on leafy vegetables within 14 days.

Use 6 tablespoons per gallon of water to control Tomato Hornworm, Tomato Fruitworm, Lace, Stink and Tarnished Plant Bug, Cucumber and Flea Beetles, Leafhoppers, Squash Bugs, Aphids, Imported Cabbage Worm, Locusts, Red Spider Mites, Armyworms on Tomatoes, Egg Plant, Pepper, Cucumber, Squash, Pumpkin and Melons. For the control of Corn earworm, apply 6 to 8 oz. as they appear. Make 3 to 4 applications at 7 day intervals. A infestation warrants re-application. Re-apply immediately after each application of this product to the tassel region of corn. After a 14 day period will seriously reduce bee population. Do not spray within 7 days of harvest. On Head Lettuce, Cabbage, Brussels Sprouts, Turnips within 7 days of harvest. On leafy vegetables, Kale, Spinach, Swiss Chard, Collards, use within 14 days of harvest. On Beets and Turnips when tops are to be used as food, use 6 tablespoons per gallon of water for control of Leafhopper, Grasshopper, Armyworm, Stink and Tarnished Plant bugs, Cutworms, Japanese Beetle and Cucumber Beetle. On Beans, Peas and Corn, Do not spray within 7 days of harvest.

FRUIT: Use 6 tablespoons per gallon of water to control of the following: Peach, Pear, Apple, Orange, Lemon, Lime, Grapes and Blueberry. Do not use within 7 days of harvest. On the Root Borer, use 6 to 8 oz. per gallon of water. Do not spray within 7 days of harvest. Do not use within 7 days of harvest.

ne. Do not apply within 7 days of harvest. For Peach
rs apply 2 or 3 sprays to trunk, from ground to scaffold
ed with moth flight.

Tablespoons per gallon of water to control Codling Moth on
pears, and Walnuts. For the control of Apple Moth, Apple
Borer, Fruit Moth, or Plum Curculio. Apply at petal fall
10 to 14 days thereafter until control is achieved.

Beetles, Leaf Rollers, Weevils on Strawberry, Boysenberry,
Blackberry. Do not use within 7 days of harvest except
berry, within 3 days.

Harmful if swallowed, inhaled, or absorbed through the
skin. Do not take internally. Skin contact may be harmful. Avoid
contact, wash hands and face before eating or smoking. Wear
gloves. Avoid contamination of food.

Product is highly toxic to bees exposed to direct treatment or
on crops. Protective information may be obtained from
Cooperative Agricultural Extension Service. Do not use on
or other uses not covered by this label. Do not leave
empty spray container in sprayer. Do not spray indoors.

Product is toxic to fish. Keep out of lakes, streams or ponds.
Do not apply where runoff is likely to occur. Do not apply when
weather conditions favor drift from areas treated. Do not contaminate
water used for cleaning of equipment, or disposal of wastes. Apply this
product only as specified on this label. Do not reuse containers.

FLAMMABLE! KEEP AWAY FROM HEAT OR OPEN FLAME.
Malathion is a cholinesterase inhibitor and can cause symptoms
similar to those caused by organic phosphate compounds. If poison-
ing occurs, call a physician immediately. Atropine is antidotal.

PHYSICIAN: Carbaryl is a moderate, reversible, Cholin-
esterase inhibitor. Atropine is antidotal.

Buyer assumes all risks of use, storage or handling of this
product not in strict accordance with directions given herewith.

No. 5719-56-AA


EPA EST. No. 5719 CA-1

CHACON CHEMICAL CORPORATION

2600 Yates Ave. • City of Commerce, CA. 90040

A Subsidiary of Leisure Enterprises, Inc.

EPA Reg. No. 5719-56



CAUTION: KEEP OUT OF REACH OF CHILDREN
See back panel for additional cautions.

NET 32 FL. OZ. (1 QUART)

CHACON

Read entire label
with label directions

Apply when insect
intervals or as needed
diluted spray per

VEGETABLES: Use
Leafhopper on Tomato
Melonworm on Cucumber
leafy vegetables. Leafy
vegetables.

Use 6 tablespoons
worm, Tomato Flea
Cucumber and Flea
Imported Cabbage
Tomatoes, Eggplant
Melons. For the cabbages
as they appear. Major
infestation warrants
of this product to period will serious
days of harvest. Cabbage, Cauliflower,
harvest. On leafy
use within 14 day
tops are to be used
for control of Leaf
Tarnished Plant Beetles
Beetles on Beans. E

FRUIT and tub
Psyllid Red Spider
30 days after bloom
Soft Brown Scale c
Do not apply during
June Beetle, San Joaquin