

PM 16 5719-56
1

00000
0
0
00000

CHACON

FRUIT AND VEGETABLE
SPRAY

ACCEPTED
APR 17 1984
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 5719-56

ACTIVE INGREDIENTS:

Malathion	6.5%
Carbaryl (1-naphthyl N-methylcarbamate)	6.5%
INERT INGREDIENTS:	87.0%

o,o-dimethyl dithiophosphate of diethyl
mercaptosuccinate.

EPA REG. NO. 5719-56

EPA EST. NO. 5719-CA-1

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for additional
precautionary statements.

NET 1 PINT
(0.473 L.)

RECEIVED

MAR 28 1984

REGISTRATION DIVISION
(TS-767)

113

FRUIT AND VEGETABLE SPRAY

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION

Keep out of reach of children. Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. Avoid contamination of food or feed.

Statement of Practical Treatment: If swallowed, do not induce vomiting, get immediate medical attention. If inhaled, remove affected person from exposure. If in eyes, flush with plenty of fresh water for at least 15 minutes, get medical attention if irritation persists. If on skin, wash affected area with soap and water.

Note to Physicians: Carbaryl and Malathion are reversible cholinesterase inhibitors. Atropine is antidotal. Do not administer I-PAM, opiates or cholinesterase inhibiting drugs.

Environmental Hazards

This pesticide is toxic to aquatic organisms. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension service.

Physical or Chemical Hazards

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Read entire label thoroughly. Apply this product only as specified on this label.

Apply when insects or damage first appears. Repeat at 7 to 10 day intervals or as necessary.

VEGETABLES: Use 3 tablespoonfuls per gallon of water. To control of Leafhoppers on Tomatoes; Pickle Worm or Melonworm on Cucumbers, Squash, Pumpkins and Melons, do not use within 3 days of harvest. Flea Beetle or Leafhoppers on Leafy Vegetables, do not use within 14 days of harvest.

Use 5 tablespoonfuls per gallon of water to control: Tomato Hornworm, Tomato Fruitworm; Lace, Stink or Tarnished Plant Bug; Cucumber or Flea Beetle, Leafhoppers, Squash Bugs, Aphids, Imported Cabbage Worm, Looper, Red Spider Mite, Armyworms on Tomatoes, Egg Plant, Peppers, Cucumbers, Squash, Pumpkins and Melons. For the control of Corn Earworm, apply to silks as soon as they appear. Make 3 to 4 applications at 3 to 5 day intervals as infestation warrants. Re-apply immediately after rain.

Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population. Do not apply within 3 days of harvest. On Head Lettuce, Carrots, Broccoli, Radish, Cabbage, Cauliflower, Brussels Sprouts, do not use within 7 days of harvest. On Leafy Vegetables, Kale, Spinach, Swiss Chard, do not use within 14 days of harvest. Do not use on Beets and Turnips when tops are to be used as food. Use 6 tablespoons per gallon of water for control of Leafhoppers, Grasshoppers, Armyworms; Stink or Tarnished Plant Bug; Caterpillars, Japanese, Flea or Cucumber Beetle on Beans, Peas and Corn. Do not use within 3 days of harvest.

FRUIT: Use 6 tablespoons per gallon of water to control Pear Psyllid, Red Spider Mite, on Apples and Pears. Do not apply until 30 days after bloom or within 3 days of harvest. Do not use on Citrus. Do not apply within 7 days of harvest. Do not apply during full bloom. Peach Leaf Curl, Plum Curculion, Pear Leafminer, Pear Leaf Scale, Pear Leaf Aphid, Nectarine, Cherry, Plum, Pear. Do not apply within 3 days of harvest. For peach tree borers apply, or spray, to trunk, to ground to scaffold limbs, timed with moth flight. Use 6 tablespoons per gallon of water to control Codling Moth on Apples, Pears, and Walnuts. For the control of Apple Moth, Apple Maggot, Oriental Fruit Moth, or Plum Curculion, apply at petal fall and every 10 to 14 days thereafter until control is achieved. Japanese Beetles, Leaf Rollers, Sawfly on Strawberry, Raspberry, Blackberry, do not use within 7 days of harvest. On Strawberry, do not use within 14 days of harvest.

STORAGE AND DISPOSAL

Storage: Store in original container in a cool, dry place away from sparks or open flame. Keep out of reach of children.
Disposal: Securely wrap original container in several layers of paper and discard in trash. Do not reuse container.

Notice: Buyer assumes all risk of use, storage and handling of this material not in strict accordance with directions given herewith.

Manufactured By
CHACON CHEMICAL CORPORATION
2600 Yates Avenue - City of Commerce, CA 90040
A GardenAmerica Company.