

**CAUTION:** Harmful if swallowed, inhaled, or absorbed through the skin. Avoid breathing of spray mist. Avoid skin contact, wash hands and face before eating or smoking. Wear clean clothing. Avoid contamination of food. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use on pets or for other uses not covered by this label. Do not leave mixed spray in sprayer. Do not spray indoors. Do not reuse containers.

This product is toxic to fish, shrimp and crab. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources.

Rinse spray equipment after using and dispose of waste and empty container by burying in non-crop land. Apply this product only as specified on this label.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME**

Malathion is a cholinesterase inhibitor and can cause symptoms similar to those caused by other organic phosphate compounds. If poisoning should occur, call a physician immediately. Atropine is antidotal.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**EPA EST. 5719-CA-1  
EPA REG. NO. 5719-59-AA**

**CHACON CHEMICAL CORPORATION  
5245 Chakemco Street • South Gate, California 90280**

A Subsidiary of Leisure Enterprises, Inc.

**CHACON**

**SCALE and  
SPIDER MITE  
CONTROL**

**ALSO CONTROLS  
APHIDS • MEALYBUGS • LEAFHOPPERS**

**ACTIVE INGREDIENTS**

Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate) ..... 6.50%  
\*Light Medium Petroleum oil ..... 81.50%

**INERT INGREDIENTS** ..... 12.00%

\*(California Dept. of Agriculture Classification of petroleum oil: Light Medium Unsulphonated Residue 92%)

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
See back panel for additional cautions.

**NET 16 FL. OZ. (1 PINT)**

**CHACON SCALE AND SPIDER MITE CONTROL**

Read entire label carefully. Use only as directed in accordance with label cautions and directions. **DO NOT** apply when temperatures are above 85°F.

**DIRECTIONS**

**SHAKE WELL BEFORE USING**

Use at the rate of 5 tablespoonfuls (2½ oz.) per gallon of water for the control of Red, Brown, Black, Purple and Cottony Cushion Scale on Citrus, Avocados, Boxwood, Camellias, Eugénias, Gardenias, Ivy (English), Oleanders, Privets. Spray thoroughly with 2 applications, 2 weeks apart. Also for Scale Insects and over-wintering forms of Spider Mites on Almonds, Apples, Plums, Pears. Apply as soon as trees begin to lose their leaves. Do not apply within 3 days of harvest. Do not apply to Avocados, Apricots, Citrus, and Peaches within 7 days of harvest.

Use at the rate of 3 tablespoonfuls (1½ oz.) per gallon of water for the control of Black, Rose, Apricot Brown and Hemispherical Scale on Boysen and other cane berries. Spray at 3-week intervals until controlled. (If berries are dormant use at the rate of 5 tablespoonfuls [2½ oz.] Also for Woolly Aphid on Apples and Pears. Also Red Spider Mites, Mealybugs, Aphids, Leafhoppers on ornamentals. Do not apply within 3 days of harvest.

Use at the rate of 2 tablespoonfuls (1 oz.) per gallon of water for the control of Red Spider Mites and Almond Mites on Almonds, Apples, Pears, Plums when in full foliage. On Apples, may injure foliage of some varieties. Use at 2-week intervals until controlled. Do not apply within 3 days of harvest. Do not apply to Avocados, Apricots, Citrus, and Peaches within 7 days of harvest.

**DO NOT** apply on Crassula, Hydrangeas, Elephant Ears and other tender foliage plants. Do not use on Coco Palm, Maiden-hair, Boston or other Ferns, African Violets, Kentia Palms.

**ACCEPTED**  
MAY 9 1975  
5719-59