

warranted, repeat applications at 7 to 10 day intervals. On apples slight russeting may occur on some varieties such as Golden Delicious. Do not spray within the following number of days before harvest: 5 days - Strawberries; 10 days - Apricots, Cherries, Nectarines, Grapes; 14 days - Apples; 21 days - Grapefruit, Lemons, Oranges, Peaches; Walnuts - Do not apply after husks open or split.

VEGETABLES: Beans, Peas, Cabbage, Cauliflower, Broccoli, Radishes, Squash, Carrots, Lettuce, Cantaloupes, Cucumbers, Sweet Corn, Onions, Peppers, Potatoes, Tomatoes; **Controls:** Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Mites, Diamondback Moths, Flea Beetles, Imported Cabbage Worms, Dipterous Leafminers, Thrips, Colorado Potato Beetles, Fall Armyworms (Tomato), Onion Thrips, Southern Armyworms (Potato and Tomato). Use 1 fl. oz. in 3 gals. of water, covers 1000 sq. ft. of garden. Do not spray within the following number of days before harvest: 1 day - Peas, Tomatoes, Sweet Corn; 3 days - Cantaloupes; 5 days - Broccoli, Cauliflower; 7 days - Beans, Cabbage, Cucumbers, Peppers, Squash; 10 days - Carrots, Lettuce, Onions, Radishes; 35 days - Potatoes.

WARNING: Keep out of reach of children. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Keep out of reach of domestic animals. Do Not Use In The Home.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

This product is toxic to fish, shrimp, crab, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective Information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse container. Destroy when empty.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

EPA REG. NO. 5719-63-AA

EPA EST. 5719-CA-1

CHACON CHEMICAL CORPORATION
2600 Yates Ave. • City of Commerce, CA. 90040
 A Subsidiary of Leisure Enterprises, Inc.

CHACON

diazinon

SPRAY

LAWN AND GARDEN

fruit and vegetable

**CONTROLS: ANTS, APHIDS, CUTWORMS, EARWIGS,
 MEALYBUGS, SOWBUGS, WHITEFLIES
 LAWN MOTHS, LEAF MINERS, LEAFHOPPERS,**

ACTIVE INGREDIENT:
 O, O-diethyl O- (2-isopropyl-6-methyl-4-pyrimidinyl)
 phosphorothioate* 25.00%

INERT INGREDIENTS 75.00%

*1 pint contains 0.24-lbs. active ingredient. **TOTAL** 100.00%

DIAZINON, ® trademark of CIBA-GEIGY

WARNING: KEEP OUT OF REACH OF CHILDREN
 See back panel for additional warnings.

NET 32 FL. OZ. (1 QUART)

CHACON DIAZINON SPRAY

Read entire label carefully before using. Use only in accordance with warnings and directions.

DIRECTIONS

Determine size of spraying job. For lawns and vegetables to be sprayed. For all other applications, apply as a thorough spray. Make applications when insects first appear, or in advance of infestation. Be sure to spray underside of leaves. Repeat applications as needed or as stated in the directions. Use contents with care.

(Note: Do not use on ferns, poinsettia, hibiscus, papaya, and orchids since injury might occur.)

LAWNS: Bent, Bermuda, Dichondra, Fescue, Merion, St. Augustine, Zoysia; **Controls:** Ants, Armyworms, Bermudagrass Thrips, Crickets, Fleas, Frit Flies, Lawn Billbugs, Lawn Moths (Cutworms), Leafhoppers, Mites (Bermuda, Clover), Sowbugs (Pillbugs), Rhodesgrass Scales; Use 1 fl. oz. in 3 gals. of water. Thoroughly wet down grass a few hours before application. Billbugs, water lawn immediately after treatment. Keep area until area has completely dried.

TREES & SHRUBS: Arborvitae, Azalea, Birch, Boxwood, Juniper, Maple, Oak, Pine, Poplar, Rhododendron, Spruce; **Controls:** Aphids, Bagworms, Leaf Miners, Mites (Clover, Cy Red, Privet, Two-Spotted Spider), Scale Crawlers (Euonymus, Pine Needle, Pit Scale, San Jose, Soft). Use of water. Apply as a thorough foliar spray. Wet both sides of branches to the dripping point, try to penetrate dense foliage.

ROSES & FLOWERS: Azaleas, Camellias, Chrysanthemum, Gladioli, Lilac, Rhododendron, Roses; **Controls:** Aphids, Scale Crawlers, Flea Beetles, Dipterous Leaf Miners, Leafhoppers, Thrips, Whiteflies. Use 1 fl. oz. in 3 gals. of water. Apply as a foliar spray. Wet the plants to the dripping point. Try to wet leaves and penetrate dense foliage.

FRUITS & NUTS: Apples, Apricots, Nectarines, Cherries, Grapes, Grapefruits, Lemons, Oranges, Peaches, Plums; **Controls:** Aphids, Apricot Mealbug, Spider Mites, Mites (Clover Spotted), Scale (Cottony-Cushion, San Jose), Codling Moth, Cherry Fruit Fly, Leafhoppers, Maggots (apples), (European Red, Pacific, Two-Spotted Spider). Use 1 fl. oz. in 3 gals. of water. Apply as a full cover spray. Start spraying after bud break in the summer. Applications should be made according to directions when aphids, mites and crawler stages of scale insects are present.

Day intervals. On apples slight
 such as Golden Delicious. Do not
 of days before harvest: 5
 cherries, Nectarines, Grapes; 14
 Lemons, Oranges, Peaches;
 or split.

auliflower, Broccoli, Radishes,
 Cucumbers, Sweet Corn, Onions,
 aphids, Cucumber beetles, Gras-
 Beetles, Mites, Diamondback
 Worms, Dipterous Leafminers,
 worms (Tomato), Onion Thrips,
). Use 1 fl. oz. in 3 gals. of water,
 y within the following number of
 Tomatoes, Sweet Corn; 3
 cauliflower; 7 days - Beans, Cab-
 lays - Carrots, Lettuce, Onions,

May be fatal if swallowed. May be
 ray mist. Avoid contact with skin,
 or handling. Avoid contamination
 household pets or humans. Keep out
 se In The Home.

NEAR HEAT OR OPEN FLAME.
), birds, and other wildlife. Birds
 e killed. Keep out of any body of
 y to occur. Do not apply when
 treated. Do not contaminate water
 wastes. Apply this product only as

ed to direct treatment or residues
 obtained from your Cooperative

empty.

Storage or handling of this material
 given herewith.

EPA EST. 5719-CA-1

CORPORATION
 mmerce, CA. 90040
 Enterprises, Inc.

CHACON

diazinon
SPRAY

LAWN AND GARDEN
fruit and vegetable

CONTROLS: ANTS, APHIDS, CUTWORMS, EARWIGS,
MEALYBUGS, SOWBUGS, WHITEFLIES
LAWN MOTHS, LEAF MINERS, LEAFHOPPERS,

ACTIVE INGREDIENT:
 O, O-diethyl O- (2-isopropyl-6-methyl-4-pyrimidinyl)
 phosphorothioate* 25.00%
INERT INGREDIENTS 75.00%
 *1 pint contains 0.24-lbs. active ingredient. **TOTAL** 100.00%
DIAZINON, ® trademark of CIBA-GEIGY

WARNING: KEEP OUT OF REACH OF CHILDREN
 See back panel for additional warnings.

NET 32 FL. OZ. (1 QUART)

CHACON DIAZINON SPRAY

Read entire label carefully before using. Use only in accordance with label warnings and directions.

DIRECTIONS

Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear, or in advance of an expected infestation. Be sure to spray underside of leaves. Repeat applications as needed or as stated in the directions. Use contents within 6 months after opening.

(Note: Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia, and orchids since injury might occur.)

LAWNS: Bent, Bermuda, Dichondra, Fescue, Merion, St. Augustine. Controls: Ants, Armyworms, Bermudagrass Thrips, Crickets, Cutworms, Earwigs, Fleas, Frit Flies, Lawn Billbugs, Lawn Moths (Sod Webworms), Leafhoppers, Mites (Bermuda, Clover), Sowbugs (Pillbugs), Springtails (Collembola), Rhodesgrass Scales; Use 1 fl. oz. in 3 gals. of water per 125 sq. ft. Thoroughly wet down grass a few hours before applying. For Lawn Billbugs, water lawn immediately after treatment. Keep children and pets off area until area has completely dried.

TREES & SHRUBS: Arborvitae, Azalea, Birch, Boxwood, Camellia, Elm, Juniper, Maple, Oak, Pine, Poplar, Rhododendron, Spruce, Willow; Controls: Aphids, Bagworms, Leaf Miners, Mites (Clover, Cyclamen, European Red, Privet, Two-Spotted Spider), Scale Crawlers (Cottony-Cushion, Euonymus, Pine Needle, Pit Scale, San Jose, Soft). Use 1 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray. Wet both sides of leaves and branches to the dripping point, try to penetrate dense foliage.

ROSES & FLOWERS: Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Rhododendron, Roses; Controls: Aphids, Euonymus Scale Crawlers, Flea Beetles, Dipterous Leaf Miners, Leafhoppers, Mealybugs, Thrips, Whiteflies. Use 1 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray. Wet the plants to the dripping point. Try to hit the underside of leaves and penetrate dense foliage.

FRUITS & NUTS: Apples, Apricots, Nectarines, Cherries, Strawberries, Grapes, Grapefruits, Lemons, Oranges, Peaches, Walnuts; Controls: Aphids, Apricot Mealbug, Spider Mites, Mites (Clover [brown], Two-Spotted), Scale (Cottony-Cushion, San Jose), Codling Moths, Fruit Leaf Roller, Cherry Fruit Fly, Leafhoppers, Maggots (apples), Mealybugs, Mites (European Red, Pacific, Two-Spotted Spider). Use 1 fl. oz. in 3 gals. of water. Apply as a full cover spray. Start spraying after blooming and again in the summer. Applications should be made according to infestation or when aphids, mites and crawler stages of scale insects are first noted. If

MAY 25 1978

Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No.

warranted, repeat applications at 7 to 10 day intervals. On apples slight russeting may occur on some varieties such as Golden Delicious. Do not spray within the following number of days before harvest: 5 days - Strawberries; 10 days - Apricots, Cherries, Nectarines, Grapes; 14 days - Apples; 21 days - Grapefruit, Lemons, Oranges, Peaches; Walnuts - Do not apply after fruits open or split.

VEGETABLES: Beans, Peas, Cabbage, Cauliflower, Broccoli, Radishes, Squash, Carrots, Lettuce, Cantaloupes, Cucumbers, Sweet Corn, Onions, Peppers, Potatoes, Tomatoes; **Controls:** Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Mites, Diamondback Moths, Flea Beetles, Imported Cabbage Worms, Dipterous Leafminers, Thrips, Colorado Potato Beetles, Fall Armyworms (Tomato), Onion Thrips, Southern Armyworms (Potato and Tomato). Use 1 fl. oz. in 3 gals. of water, covers 1000 sq. ft. of garden. Do not spray within the following number of days before harvest: 1 day - Peas, Tomatoes, Sweet Corn; 3 days - Cantaloupes; 5 days - Broccoli, Cauliflower; 7 days - Beans, Cabbage, Cucumbers, Peppers, Squash; 10 days - Carrots, Lettuce, Onions, Radishes; 35 days - Potatoes.

WARNING: Keep out of reach of children. May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Keep out of reach of domestic animals. Do Not Use In The Home.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

This product is toxic to fish, shrimp, crab, birds, and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse container. Destroy when empty.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

EPA REG. NO. 5719-63-AA EPA EST. 5719-CA-1

CHACON CHEMICAL CORPORATION

2600 Yates Ave. • City of Commerce, CA. 90040

A Subsidiary of Leisure Enterprises, Inc.

CHACON

diazinon
SPRAY

LAWN AND GARDEN
fruit and vegetable

CONTROLS ANTS APHIDS CUTWORMS EARWIGS
MEALYBUGS, SOWBUGS, WHITE FLIES
LAWN MOTHS, LEAFMINERS, LEAFHOPPERS

ACTIVE INGREDIENT:
O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)
phosphorothioate 25.00%
INERT INGREDIENTS 75.00%
*1 pint contains 0.24-lbs. active ingredient. **TOTAL** 100.00%
DIAZINON, ® trademark of CIBA-GEIGY

WARNING: KEEP OUT OF REACH OF CHILDREN
See back panel for additional warnings.

NET 32 FL. OZ. (1 QUART)

CHACON DIAZINON SPRAY

Read entire label carefully before using. Use only in accordance with warnings and directions.

DIRECTIONS

Determine size of spraying job. For lawns and vegetables, measure to be sprayed. For all other applications, apply as a thorough cover. Make applications when insects first appear, or in advance of an infestation. Be sure to spray underside of leaves. Repeat applications needed or as stated in the directions. Use contents within 6 months opening.

(Note: Do not use on ferns, poinsettia, hibiscus, papaya, pine, and orchids since injury might occur.)

LAWNS: Bent, Bermuda, Dichondra, Fescue, Merion, St. Augustine; **Controls:** Ants, Armyworms, Bermudagrass Thrips, Crickets, Grasshoppers, Fleas, Frit Flies, Lawn Billbugs, Lawn Moths (and Webworms), Leafhoppers, Mites (Bermuda, Clover), Sowbugs (Flatbugs), Spring (Collembola), Rhodgrass Scales; Use 1 fl. oz. in 3 gals. of water per sq. ft. Thoroughly wet down grass a few hours before applying. For Billbugs, water lawn immediately after treatment. Keep children and pet area until area has completely dried.

TREES & SHRUBS: Arborvitae, Azalea, Birch, Boxwood, Camellia, Juniper, Maple, Oak, Pine, Poplar, Rhododendron, Spruce, Willow; **Controls:** Aphids, Bagworms, Leaf Miners, Mites (Clover, Cyclamen, European Red, Privet, Two-Spotted Spider), Scale Crawlers (Cottony-Cup Euonymus, Pine Needle, Pin Scale, San Jose, Soft). Use 1 fl. oz. in 3 of water. Apply as a thorough foliar spray. Wet both sides of leaf branches to the dripping point, try to penetrate dense foliage.

ROSES & FLOWERS: Azaleas, Camellias, Chrysanthemums, Carna, Gladioli, Lilac, Rhododendron, Roses; **Controls:** Aphids, Euonymus Crawlers, Flea Beetles, Dipterous Leaf Miners, Leafhoppers, Mealy Thrips, Whiteflies. Use 1 fl. oz. in 3 gals. of water. Apply as a thorough foliar spray. Wet the plants to the dripping point. Try to hit the underside leaves and penetrate dense foliage.

FRUITS & NUTS: Apples, Apricots, Nectarines, Cherries, Strawberries, Grapes, Grapefruit, Lemons, Oranges, Peaches, Walnuts; **Controls:** Aphids, Apricot Mealybug, Spider Mites, Mites (Clover (brown), Spotted), Scale (Cottony-Cushion, San Jose), Codling Moths, Fruit Fly, Cherry Fruit Fly, Leafhoppers, Maggots (apples), Mealybugs (European Red, Pacific, Two-Spotted Spider). Use 1 fl. oz. in 3 of water. Apply as a full cover spray. Start spraying after blooming and in the summer. Applications should be made according to infestation when aphids, mites and crawler stages of scale insects are first noticed.