

ACC
 JUN 6 1969
 Under the
 Fungicide
 as used in the
 in the
 5719-69

5719-69

VEGETABLES for the control of red spider, tomato russet, Pacific, Tropical mites on beans, peppers, squash (winter and summer), tomatoes and all melons. Do not use on beans within 21 days of harvest and 2 days of harvest on other crops listed. Do not use on eggplant.

STRAWBERRIES for control of six-spotted, strawberry and two-spotted mites. Use at the rate of 2 teaspoonfuls per gallon of water. Apply when mites appear. Repeat as necessary. Do not apply within 2 days of harvest.

FRUITS AND NUTS: Use at the rate of 2 teaspoonfuls per gallon of water. Apples, Crabapples, Pears, Quince to control the following mites: Clover, European Red Mite, Apple Mite, McDaniel, Pacific, Schoene, Red Spider, Willamette and Yellow (Carpini) Mites. Do not apply within 7 days of harvest. Apricots, Cherries, Figs, Nectarines, Peaches, Plums, Prunes, to control mites listed above. Apply at petal fall or when mites appear. Wait 30 days to repeat treatment if necessary. Do not spray within 7 days of harvest. Do not use on Avocados, Grapes, Chestnuts, Filberts, Pecans. Walnuts to control Brown almond (clover), European Red, McDaniel, Pacific, two spotted Willamette and yellow mites. Apply just after bloom or whenever mites appear and repeat if necessary. Do not apply within 14 days of harvest. **Clover Mites in Building:** Use at the rate of 2 tablespoonfuls per gallon of water. Spray thoroughly outside walls, foundations, windowsills, plants and lawns at base of infested buildings.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

EPA Est. 5719-CA-1 EPA Reg. No. 5719 69
CHACON CHEMICAL CORPORATION
 2600 Yates Avenue - City of Commerce, California 90040
 A Subsidiary of Leisure Enterprises, Inc.

CHACON

KELTHANE
MITE CONTROL

**RED SPIDER MITE, CLOVER MITE, BERMUDA MITE,
 TOMATO RUSSET MITE, STRAWBERRY MITE,
 BROWN MITE**

| | | |
|-------------------|--|---------|
| ACTIVE INGREDIENT | 1,1 Bis(chlorophenyl) 2,2,2 trichloroethanol | 18.50% |
| INERT INGREDIENTS | | 81.50% |
| | | 100.00% |

*Equivalent to 1.6 lbs. active ingredient per gallon.
 U.S. Patents 2,812,166 & 2,812,167

KEEP OUT OF REACH OF CHILDREN
CAUTION

In case of contact, wash skin with soap and water. If eyes flush with water for 15 minutes and get medical attention.
 See side panel for additional precautionary statements.

NET 16 FL. OZ. (1 PINT)

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION
 Keep out of reach of children. Avoid skin contact. Wash thoroughly after using. Avoid inhalation of mist. Harmful if swallowed. If taken internally, get medical attention.

ENVIRONMENTAL HAZARD

This pesticide is toxic to fish. Use with care when applying in areas adjacent to any body of water. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated.

PHYSICAL - CHEMICAL HAZARDS

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME.

DIRECTIONS

Do not store near feed or food stuffs.

DISPOSAL

WRAP CONTAINER AND PUT IN TRASH COLLECTION.

Read entire label carefully. Use only in accordance with label cautions and directions.

ORNAMENTALS, TREES, NURSERY STOCK: Use at the rate of 2 teaspoonfuls per gallon of water for the control of Red Spider Mite (two-spotted). Apply when mites appear. Repeat as necessary.

LAWNS for the control of Bermuda, brown and clover mites. Use at the rate of 2 tablespoonfuls (1 fl. oz.) per gallon of water per 400 square feet of lawn. Make application when mites appear. Spray the entire lawn area with a thorough spray and repeat in 10 to 14 days if necessary.