

disinfectant, sanitizer and algicide

DANGER: CAUSES SEVERE EYE DAMAGE AND SKIN IRRITATION.

Do not get in eyes, on skin or clothing. Harmful if swallowed. Avoid contamination of food. Wear goggles or face shield when handling. Remove and wash contaminated clothing before re-use.

FIRST AID: FLUSH EYES OR SKIN WITH PLENTY OF WATER FOR AT LEAST 15 MINUTES. FOR EYES CALL A PHYSICIAN. REMOVE AND WASH CONTAMINATED CLOTHING BEFORE RINSE. IF SWALLOWED DO NOT INDUCE VOMITING. DRINK LARGE QUANTITIES OF FLUIDS AND CALL A PHYSICIAN IMMEDIATELY. RINSE CONTAINER AND DISCARD IT.

THIS PRODUCT IS TOXIC TO FISH. TREATED EFFLUENT WATERS MAY BE HARMFUL TO FISH AND OTHER AQUATIC LIFE.

KEEP OUT OF THE REACH OF CHILDREN

FOR CONTROL OF SLIME & ALGAE: In Industrial Water Treatment and Cooling Towers add Initial Dose of 1 - 2 gallons of D-TROL to 5,000 gallons of water. Use a Periodic Dose of 1 - 3 quarts of D-TROL to 5,000 gallons of water every 7 days or sooner should visible growth appear. For best results, rid system of organic debris if possible. If soap or anionic detergent is used, rinse thoroughly before charging with D-TROL.

FOR SANITIZING EQUIPMENT IN MEAT AND FOULTRY PLANTS: Clean equipment with a suitable detergent and rinse. Sanitize with 1/4 ounce of D-TROL in 1 gallon of water (200 ppm Quat). Rinse with potable water after sanitizing. Listed as a Code F, sanitizer and Code N, compound for treating closed systems in the "List of Chemical Compounds" authorized for use under USDA Poultry, Meat, Rabbit and Egg Product Inspection Program.

FOR FINAL SANITIZING RINSING IN FOOD PLANTS: Clean equipment with a suitable detergent and rinse. For final sanitizing rinsing use 1/4 ounce of D-TROL (200 ppm Quat) per 1 gallon of water. D-TROL complies with Paragraph 121.2547, Sanitizing Solution, Part 121, Food Additive Law for final sanitizing rinsing of food processing equipment and on food contact surfaces in bars and restaurants. No rinse required in dairies unless specified by area health authorities. D-TROL fulfills criteria of Appendix F as revised in the Grade "A" Pasteurized Milk Ordinance and Code (1965) in water up to 750 ppm hardness when tested by Chamber Method.

FOR DISINFECTING IN HOSPITALS, REST HOMES AND VETERINARY OPERATIONS: Clean equipment with a suitable detergent and rinse. For disinfecting use 3/4 ounces of D-TROL per 1 gallon of water (600 ppm Quat) on hard surfaces. D-TROL at this concentration passes the A.O.A.C. Use Dilution Confirmation Test against Pseudomonas aeruginosa (ATCC No. 15442), Salmonella choleraesuis (ATCC No. 10708) and Staphylococcus aureus (ATCC No. 6538).

FOR USE AS ALGICIDE IN SWIMING POOL:


Initial Dose: 1 gallon of D-TROL to 50,000 gallons of water.

Maintenance Dose: 1 quart of D-TROL to 50,000 gallons of water.

Use Initial Dose when filling with fresh water. If pool has visible algae, the Initial Dose should be repeated after algae debris are removed from pool...

Maintenance Dose added every 5 to 7 days. Add 1 quart of D-TROL for each 10,000 gallons of fresh make up water per week, continue to maintain ppm of chlorine and pH of pool water specified by public health authorities.

For Winter Maintenance add 1 gallon of D-TROL to 50,000 gallons of water held in pool. Repeat if algae show growth.



ממחתום פוודואוומעדה מדוגדוער מנונסרטי מואמואוענול מנוס מימיטי