

100-5736-56

1/2

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

JAN 27 1990

Walden Chemical, Inc.
2630 East Kemper Road
Charonville, OH 45211

Attention: Joe L. Jones

Dear Mr. Jones:

Reference is made to your letter of January 17, 1990, regarding the registration of your product, "Walden Chemical, Inc. Antimicrobial Concentrate".

Place EDTA and isopropyl alcohol in the inert ingredients statement of your label. Also add it accordingly.

WJ

William G. Francis
Active Product Manager (30)
Antimicrobial Program Branch
Registration Division (E75050)

Enclosure

BEST AVAILABLE COPY

58622:1:A-4:Pringle:LK-20:KENCO:01/17/90:02/14/90:ds:sw:vo:dd:dq

CONCURRENCES

SYMBOL	H-750K						
SURNAME	Pringle						
DATE	1-19						

ONE GALLON (3.785 LITERS)
U.S. MEASURE
NOT SOLD FOR RESIDENTIAL USE
DuBOIS CHEMICALS, INC.

FUNGISTATIC BARRICADE

fungistatic barrier agent

ACTIVE INGREDIENTS	19.20%
Tetrasodium ethylene diamine tetracetate	0.40%
o-Benzyl-p-chlorophenol	3.60%
Isopropyl alcohol	11.20%
INERT INGREDIENTS	84.80%
	100.00%

KEEP OUT OF REACH OF CHILDREN
DANGER:

SEE PRECAUTIONARY STATEMENT
ON SIDE OF CONTAINER

All surfactants used in this product are biodegradable

This product does not contain phosphorus
compounds. Composition complies with
EPA testing procedures for a fungistatic film

EPA Registration No. 5736-56
EPA Est. No. 5736 OH-1

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

The surfaces should be clean and reasonably dry. FUNGISTATIC BARRICADE is best applied with a pressure spray tank with a fine nozzle and a 15 to 20 psi tank pressure. Apply a continuous film, saturating the surface from front to back. It also can be brushed for application on equipment or dipped. The amount should be the minimum to accomplish the desired effect. FUNGISTATIC BARRICADE should be diluted 1:1 for application but if desired it can be applied undiluted. Naturally, any foodstuff in the area must be removed during spraying application to prevent foodstuff contamination. Ventilation must be maintained to prevent odor transfer.

FUNGISTATIC BARRICADE can be diluted 1:1 with water for application to troublesome mold growth areas in food and beverage plants, primarily walls and ceilings. FUNGISTATIC BARRICADE should be applied only to non-food contact areas. When dried to a firm coating (usually 30 to 60 minutes), FUNGISTATIC BARRICADE will inhibit the growth of mold and fungus and also act as a soil release film. In high humidity areas, a longer drying time may be needed. Adequate ventilation must be maintained during spraying to remove volatile components of product.

The coverage for FUNGISTATIC BARRICADE when diluted 1:1 is 600 square feet per gallon.

DuBois Chemicals, Inc. • DuBois Tower
Cincinnati, Ohio 45202

For additional customer service, call 800-443-8311. For technical assistance, call 513-752-1111. To contact your representative, call 513-752-1111. For more information, call 513-752-1111.



Product No. 22452

FUNGISTATIC BARRICADE

fungistatic barrier agent

Precautionary Statements

Hazards to Humans and Domestic Animals.

KEEP OUT OF REACH OF CHILDREN.

DANGER: Flammable. Defats skin and irritates eyes, nose, throat, and on clothing. Protect eyes and skin when handling. Wash thoroughly after use. Avoid contamination of food.

STATEMENT OF PRACTICAL TREATMENT: In case of contact, immediately flush eyes with plenty of water for at least 15 minutes. For eye contact, remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of water. Avoid alcohol. **PHYSICIAN IMMEDIATELY.**

ENVIRONMENTAL HAZARDS: Do not discharge into lakes, streams, or public waters unless in accordance with NPDES permit. For guidance, contact the Regional Office of the EPA.

PHYSICAL HAZARDS: Do not use, spill or store near heat or open flame. **STORAGE AND DISPOSAL**

- Do not contaminate water, food, or feed by storage or disposal.
- Open dumping is prohibited.
- Do not reuse empty container.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. Wastes cannot be disposed of by use according to label instructions. State Pesticide or Environmental Control Agency, or the Hazardous Waste Unit at your EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling, or puncture and dispose of in a sanitary landfill, or by other means approved by state and local authorities, by burning, if burned, state and local authorities.

GENERAL: Consult federal, state or local disposal authorities for a complete list of procedures such as limited open burning.