

**Precautionary Statements
Hazards to Humans & Domestic Animals
KEEP OUT OF REACH OF CHILDREN.
CAUTION**

Harmful if swallowed. Do not apply while food processing is underway. Except in federally inspected meat and poultry plants, when applied as a space spray with care and in accordance with the directions and precautions given above, food processing operations may continue. Foods should be removed or covered before treatments. All food processing surfaces should be covered during treatments or thoroughly washed and rinsed with potable water before reuse. Stored food products which are infested should be fumigated or treated by other effective methods using an approved product intended for that purpose.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL

Pesticide spray mixture or rinsate that cannot be used according to label instructions must be disposed of according to Federal, State or local procedures for the Resource Conservation and Recovery Act.

CONTAINER DISPOSAL

Triple rinse (or equivalent) and offer for recycling, reconditioning or dispose of in a sanitary landfill or by other State and local procedures.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herein.

DuTOXIDE 550

PYRETHRIN SPACE SPRAY


DuTOXIDE 550 is a stable synergized pyrethrin Micro Emulsion—Leaves No Oily or Objectionable Residue. Designed for Use in Bakeries, Beverage Plants, Canneries, Food Processing Plants, Truck Trailers, Warehouses, Institutions, Restaurants, Barns, Storage Rooms, Mushroom Production and Processing, and Outdoors for Mosquito Control.

Apply with Conventional Mechanical or Compressed Air Equipment Designed and Adjusted to Deliver a True Aerosol Mist.

ACTIVE INGREDIENTS

- Pyrethrins
- Pyperonyl Butoxide
- Permethrin

INERT INGREDIENTS



- 05%
- 50%
- 20%
- 9.25%
- 100%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Harmful if swallowed. Avoid contamination of feed and foodstuffs. See below and side panels for additional precautionary statements.

Net contents of each container
EPA Registration No. 5736-75 AA • EPA No. 5736-001

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SPACE APPLICATIONS

For rapid control of: Flies, Fruit Flies, Gnats, Mosquitoes, Flies, Wasps, Hornets, Angoumois Grain Moths, Mediterranean Moths, Indian Meal Moths, and Tobacco Moths.

Close windows and doors and shut off ventilating fans. Apply DuTOXIDE 550 with a good machine adjusted to deliver a fine mist of fog. Use a fogger or vaporizer adjusted to deliver a fine mist (no droplets over 50 microns in diameter and 80% under 10 microns). Apply at the rate of 1 ounce per 1000 cubic feet. Direct the spray toward the ceiling and upper corners and behind obstructions. Keep the area closed for at least 30 minutes. Vacate the treated area and ventilate before reoccupancy. Repeat treatment as necessary. A combination of spot spraying and fogging is recommended.

IN HORSE BARN AND STORAGE ROOMS

To Control House Flies, Stable Flies and Horn Flies: Apply DuTOXIDE 550 with 4 parts of water. Apply as a fine mist to the horses. Use a fogger or mist generator to direct the mist toward the center of the area where animals are kept. Apply 1 ounce per 1000 cubic feet uniformly throughout the area sprayed. Do not ventilate the area for at least 30 minutes after spray has been applied. Repeat daily in the morning or as required.

In Mushroom Production and Processing: To Control Fungus Gnats: In growing houses, apply DuTOXIDE 550 at the rate of 1 ounce per 1000 cubic feet of space. Fogger should be placed at least 6 feet away from beds and directed slightly upward. Ventilating fans should be left in operation, but for best results, fans should be turned off in packing and canning. Apply at the rate of 1 ounce per 1000 cubic feet of space.

SURFACE SPRAYING

To Control Roaches, Ants, Silverfish, Spiders, Mites, Cheese Mites, Granary Weevils, Rice Weevils, Flour Beetles, Rust Red Flour Beetles, Sawtoothed Flour Beetles, Spider Beetles, Cigarette Beetles, Drug



DuBOIS CHEMICALS, A DIVISION OF CHEMED CORPORATION


CONTACT YOUR DuBOIS REPRESENTATIVE FOR ADDITIONAL USE DIRECTIONS AND MATERIAL SAFETY DATA SHEET.

DuTOXIDE 550

PERMETHRIN SPACE SPRAY

synergized pyrethrin Micro Emulsion—Leaves No Residue. Designed for Use in Bakeries, Beverage Plants, Food Processing Plants, Truck Trailers, Warehouses, Institutions, Greenhouses, Mushroom Production and Processing, and Storage Rooms.

Use with Mechanical or Compressed Air Equipment Designed for True Aerosol or Mist.



05
50
200
925
1000

OUT OF REACH OF CHILDREN CAUTION

Prevents contamination of feed and foodstuffs. See below for additional precautionary statements.

Labels shown elsewhere on this container.
EPA Reg. No. 57.36-75-AA • EPA Reg. No. 57.36-75-01

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

SPACE APPLICATIONS

For rapid control of: Flies, Fruit Flies, Gnats, Mosquitoes, Skipper Flies, Wasps, Hornets, Angoumois Grain Moths, Mediterranean Flour Moths, Indian Meal Moths, and Tobacco Moths.

Close windows and doors and shut off ventilating systems. Apply DuTOXIDE 550 with a good machine adjusted to deliver a fine mist (1-5 microns). Use a fogger or vaporizer adjusted to deliver an aerosol spray (no droplets over 50 microns in diameter and 80% less than 30 microns). Apply at the rate of 1 ounce per 1000 cubic feet of space. Direct the spray toward the ceiling and upper corners of the area and behind obstructions. Keep the area closed for at least 15 minutes. Vacate the treated area and ventilate before reoccupying. Repeat treatment as necessary. An approved insecticide surface treatment is recommended.

IN HORSE BARN AND STORAGE ROOMS

To Control House Flies, Stable Flies and Horn Flies: Dilute 1 part of DuTOXIDE 550 with 4 parts of water. Apply as a fine mist over the animals and to the center of the area where animals are housed. Apply 1 ounce per 1000 cubic feet uniformly throughout the entire area sprayed. Do not ventilate the area for at least 10 minutes after spray has been applied. Repeat daily in the morning and afternoon if necessary.

In Mushroom Production and Processing: To control Mushroom Flies or Fungus Gnats in growing houses, apply at a rate of 1 ounce per 1000 cubic feet of space. Fogger should be operated 8 to 10 feet away from beds and directed slightly upward over them. Circulating fans should be left in operation, but for best results, refrigeration should be turned off in packing and canning plants. Use at a rate of 1 ounce per 1000 cubic feet of space.

SURFACE SPRAYING

To Control Roaches, Ants, Silverfish, Spiders, Crickets, Clover Mites, Cheese Mites, Granary Weevils, Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Sawtoothed Grain Beetles, Spider Beetles, Cigarette Beetles, Drugstore Beetles,

Mealworm, Grain Mites, and Caddis: Use a good sprayer adjusted to deliver a coarse wet spray. Direct the spray into hiding places, cracks and crevices, under pallets, around containers of stored foods, around the base of machinery, behind shelves and drawers. For silverfish, spray bookcases. For ants, spray trails, nests and points of entry. If surface application only is to be used, spray floors, walls and other surfaces applying at a rate of 1 gallon to 750 square feet of surface.

To Control Carpet Beetles: Spray edges of carpeting and under carpeting and rugs. Make a localized application to floor and baseboards. Spray directly into cracks, closets, and infested areas of shelving. Repeat treatment as necessary.

To Control Bedbugs: Spray mattresses lightly, particularly around tufts and seams. Take beds apart and spray in all joints. Treat baseboards, moldings and floors. Repeat treatment as needed.

To Control Fleas and Brown Dog Ticks: Thoroughly spray infested areas, pet beds, resting quarters, nearby cracks and crevices along and behind baseboards, moldings, window and door frames and localized areas of floor and floor covering. Fresh bedding should be placed in animal quarters following treatment. Concurrent treatment of animals with an approved insecticide is recommended. Repeat treatment as needed.

To Control Clothes Moths: Remove any infested articles from storage, brush thoroughly and air for several hours in sunlight if possible. Apply spray liberally to empty chests, closets, bureaus, and other storages directing the spray into cracks, joints, and crevices. After airing infested articles may be treated lightly keeping nozzle at least three feet from fabric to avoid staining. Store clothes in moth tight containers or repeat treatments monthly.

OUTDOOR APPLICATIONS

Adult Mosquito Control: Apply as a fine mist to thoroughly treat the entire area. For best results, treat when the wind velocity does not exceed 5 miles per hour. Treat shrubbery or vegetation where mosquitoes may rest, low marsh areas, stagnant pools or ponds and shorelines. Apply at a rate of 10 fluid ounces per acre (0.2 fluid ounce per 1000 square feet).

Using truck-mounted conventional misting equipment traveling at 5 m.p.h. to treat a 300 foot swath, calibrate equipment to deliver 0.23 fluid ounces per minutes.

SOLELY FOR NON-FEDERAL USE • NOT FOR SALE TO EMPLOYEES OR BY SERVICE PERSONS

DuTOXIDE 550, A DIVISION OF CHEMED CORPORATION • CINCINNATI, OHIO 45202 • U.S.A.

CONTACT YOUR DuBOIS REPRESENTATIVE FOR ADDITIONAL USE DIRECTIONS AND MATERIAL SAFETY DATA SHEET