

prentox

CO-RAX PELLETED BAIT

A TOX-HID* COATED WARFARIN PELLETED BAIT
Kills Norway Rats, Roof Rats and House Mice.
For Indoor Use Only

Each particle of the Warfarin in this product has been specially coated with a material that reduces detection of the Warfarin by the rodents at the time the bait is consumed. The coating may help assure greater acceptance of the bait in cases where rodent awareness of uncoated Warfarin is responsible for low bait acceptability.

ACTIVE INGREDIENT:

Warfarin (3-alpha-Acetylbenzyl)-
4-hydroxycoumarin) 0.025%

INERT INGREDIENTS: 99.975%
TOTAL: 100.00%

* TOX-HID - Wisconsin Alumni Research Foundation - Patent Pending.
TOX-HID - Trademark of Wisconsin Alumni Research Foundation.

CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

BEST DOCUMENT AVAILABLE



Manufactured by

prentiss
Drug & Chemical Co., Inc.

Plant: 336 Wilson Avenue
Newark, N.J. 07105
Office: 363 7th Avenue
New York, N.Y. 10001

PRECAUTIONARY STATEMENTS:

HAZARD TO HUMANS & DOMESTIC ANIMALS:

CAUTION: Keep away from humans, domestic animals and pets. If swallowed by humans, domestic animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of Vitamin K₁ combined with blood transfusions are indicated, as in the case of hemorrhage caused by overdoses of bishydroxycoumarin.

ENVIRONMENTAL HAZARD:
Keep Out of Lakes,
Streams or Ponds.

DIRECTIONS FOR USE General Classification

It is a violation of federal law to use this product in a manner inconsistent with its labeling

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS & MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. Repeat treatment when infestation recurs. Where continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

USE RESTRICTIONS - For control of Norway rats, Roof rats and House mice in homes, industrial and agricultural buildings, and similar man-made structures. For indoor use only. Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS - After removing as much food, water, and harborage as possible, determine areas where rats or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in location where rodents or their signs have been observed.

STORAGE AND DISPOSAL

DO NOT CONTAMINATE WATER, FOOD OR FEED BY STORAGE OR DISPOSAL. OPEN BURNING AND DUMPING IS PROHIBITED. DO NOT REUSE EMPTY CONTAINER.

PESTICIDE DISPOSAL: Pesticide that cannot be used, should be disposed of in a landfill approved for pesticides or buried in a safe place, away from water supplies.

CONTAINER DISPOSAL: Triple rinse (or equivalent) and dispose of in an incinerator or landfill approved for pesticide containers or bury in a safe place. Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning.

EPA Reg. No. 655-539

NET CONTENTS

ACCEPTED

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 655-539

Proper timing application rate, and repeated spraying are necessary for fruit insect control. Always consult your local Agricultural Extension Specialist for fruit tree spray schedule.

DIRECTIONS FOR USE

Mix one pint Fruit and Vegetable Spray in 40 gallons of water or one ounce (2 tablespoons) in 3 gallons of water to spray fruit trees.

NOTE: Do not spray apples, pears, grapes, or cherries within 2 days of harvest. Do not spray peaches, quinces, plums, or apricots within 7 days of harvest. Do not spray more than 200 gallons per acre. Do not spray when bees are present. Do not spray when birds are present. Do not spray when bees are present. Do not spray when birds are present.

APPLES AND PEARS: Coat with spray when blossoms have fallen and repeat at 10-day intervals. Usually 4 or 5 sprayings are necessary to protect fruit from bees. Spray when cloudy and green apples (red-dipped half red) and half-ripe apples (red-dipped half red) are present. Repeat at 10-day intervals until parts are controlled.

PEACHES: Spray when half-ripe or nearly ripe. Repeat at 10-day intervals as necessary. Do not spray more than 200 gallons per acre. Do not spray when bees are present. Do not spray when birds are present.

GRAPES: Spray when half-ripe or nearly ripe. Repeat at 10-day intervals as necessary. Do not spray more than 200 gallons per acre. Do not spray when bees are present. Do not spray when birds are present.

PLUMS: Do not spray peaches, quinces, plums, or apricots within 7 days of harvest. Do not spray more than 200 gallons per acre. Do not spray when bees are present. Do not spray when birds are present.

APRICOTS: Do not spray peaches, quinces, plums, or apricots within 7 days of harvest. Do not spray more than 200 gallons per acre. Do not spray when bees are present. Do not spray when birds are present.

CHERRIES: Do not spray peaches, quinces, plums, or apricots within 7 days of harvest. Do not spray more than 200 gallons per acre. Do not spray when bees are present. Do not spray when birds are present.

QUINCES: Do not spray peaches, quinces, plums, or apricots within 7 days of harvest. Do not spray more than 200 gallons per acre. Do not spray when bees are present. Do not spray when birds are present.

PLUMS: Do not spray peaches, quinces, plums, or apricots within 7 days of harvest. Do not spray more than 200 gallons per acre. Do not spray when bees are present. Do not spray when birds are present.

APRICOTS: Do not spray peaches, quinces, plums, or apricots within 7 days of harvest. Do not spray more than 200 gallons per acre. Do not spray when bees are present. Do not spray when birds are present.

CHERRIES: Do not spray peaches, quinces, plums, or apricots within 7 days of harvest. Do not spray more than 200 gallons per acre. Do not spray when bees are present. Do not spray when birds are present.

QUINCES: Do not spray peaches, quinces, plums, or apricots within 7 days of harvest. Do not spray more than 200 gallons per acre. Do not spray when bees are present. Do not spray when birds are present.



RECEIPT

MAY 19 1978

869-152

Green Light Company
 2501 West 10th Street
 San Antonio, Texas 78217

Green Light Company
 2501 West 10th Street
 San Antonio, Texas 78217

KEEP OUT OF THE REACH OF CHILDREN

STATEMENT OF PRACTICAL TREATMENT
 If swallowed give immediate first aid. If in eyes, flush eyes with water for 15 minutes. Get medical attention.

Manufactured by GREEN LIGHT COMPANY, San Antonio, Texas 78217

Do not spray when bees are present. Do not spray when birds are present.