

UNDER THE FEDERAL INSECTICIDE
ACT, 1947
EPA REG. NO. 105-100000-000000
105-100000-000000

TERR-O-GAS 70

PREPLANT SOIL FUMIGANT

ACTIVE INGREDIENTS:

	by Wt.
Methyl Bromide	70%
Chloropicrin	30%
Total	100%

For control of Nematodes, Wireworms, Weed Seed, Soil Borne Insects and Fungus Diseases.



DANGER



KEEP OUT OF REACH OF CHILDREN

POISON

DO NOT INHALE VAPORS • SEE ANTIDOTE & WARNINGS ON SIDE PANEL

Manufactured by

GREAT LAKES CHEMICAL CORPORATION

P. O. BOX 2200, WEST LAFAYETTE, INDIANA 47906

USDA Reg. No. 5785-19



POISON



DANGER

**MAY BURN SKIN AND EYES
POISONOUS IF INHALED**

Do Not Breathe Vapor

Do Not Get in Eyes, on Skin or Clothing

In case of contact, immediately flush skin or eyes with copious amounts of water at least 15 minutes and get medical attention if eye contact. If liquid has spilled on clothing, remove from immediately and air the clothing. Do not re-use clothing until it has been thoroughly cleaned.

POISON


SEND FOR DOCTOR IN CASE OF ACCIDENT

ANTIDOTE: In case of contact, flush skin or eyes with copious amounts of water at least 15 minutes and get medical attention if eye contact. If liquid has spilled on clothing, remove from immediately and air the clothing. Do not re-use clothing until it has been thoroughly cleaned.

NOTICE

Following soil fumigation, undesirable concentrations of chloropicrin may drift to nearby areas. If this occurs, immediately cover treated area with a plastic tarpaulin. Leave tarpaulin on overnight and remove during the daytime. Recover areas if eye irritation indicates that undesirable concentrations are continuing. These concentrations are most likely to occur in the evening or at night when the air is most likely to be static in movement. Do not make application of Terr-O-Gas 70 when there is little or no air movement or there is an inversion condition in the atmosphere.

NOTE CAREFULLY: Following fumigation, the level of soluble salts and ammonia nitrogen may be raised. This is most likely to occur when heavy rates of fumigant and fertilizers are applied to soils that are either acid, wet, cold, or high in organic matter. Fertilizers containing ammonium salts should not be used. Apply only fertilizers containing nitrates until after the crop is well established and the soil temperature is above 65° F. in order to avoid ammonia injury, nitrate starvation, or both. After fumigation fertilize as indicated by soil test to protect against injury to plant roots. It is best that highly acid soils be limed before application to stimulate nitrification.



WARNING

NOTE. Many pesticidal chemicals are poisonous and may leave a toxic residue on the plants to which they are applied. The U. S. Food and Drug Administration has established minimum amounts of such pesticidal chemicals that may remain on raw agricultural products at harvest, and it is the user's responsibility to see that there is no residue on such crops at harvest in excess of these amounts. The "Suggestions for Use" above, are based on the best available information, and if followed carefully should not leave excessive residues at harvest. However, Great Lakes Chemical assumes no responsibility as to their accuracy nor for any loss due to excessive residues.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.

Empty containers should be returned to Great Lakes Chemical Corp.

TERR-O GAS FUMIGATION DIRECTIONS RECOMMENDED APPLICATION RATES

Strawberries 225 to 325
lbs/Acre

Expose to fumigation for 48 hrs. Aerate 2 weeks before setting transplants in treated areas.

Tomatoes 225 to 325
lbs/Acre

Expose to fumigation for 48 hrs. Aerate 2 weeks before setting transplants in treated areas.

Apply with chisels spaced 12 inches apart to a depth of 8 inches. Seal fumigant with a drag or cultipacker immediately behind chisels. Cover with a gas-proof tarpaulin. Remove tarpaulin after recommended fumigation period and aerate soil for two weeks before planting. If odors persist disc or plow the soil to assist aeration.

Do not apply Terr-O-Gas 70 when there is little or no air movement.

Do not apply near buildings inhabited by humans or livestock or in areas adjacent to fields where valuable crops are growing.

NOTE: Vapors of Terr-O-Gas 70 are extremely irritating to the eyes. When applying or transferring to other containers, provide good ventilation. In the presence of discomforting concentrations of Terr-O-Gas 70 vapor wear a full face mask with an approved Black Canister for acid gas and organic vapors.

GREAT LAKES CHEMICAL CORPORATION
P. O. BOX 2200,
WEST LAFAYETTE, INDIANA 47906
19523 SOUTH SUSANA ROAD
COMPTON, CALIFORNIA 90221

MADE IN U.S.A.