

WARNING
KEEP AWAY from FEED or FOOD PRODUCTS
POISON
CAUTION—DO NOT DROP
IF BREAKING DON'T BREATHE FUMES
TOUCH CONTENTS SWALLOW

This is to certify that the contents of this package are properly described by name and are packed and marked and are in proper condition for transportation according to the Regulations prescribed by the Interstate Commerce Commission.
Great Lakes Chemical Corporation
El Dorado, Arkansas

Shipping name required hereon for shipments by EXPRESS

WARNING

NOTE: Many pesticidal chemicals are poisonous and may leave a toxic residue on the plants to which they are applied. The U. S. Food and Drug Administration has established minimum amounts of such pesticidal chemicals that may remain on raw agricultural products at harvest, and it is the user's responsibility to see that there is no residue on such crops at harvest in excess of these amounts. The "Suggestions for Use" above, are based on the best available information, and if followed carefully should not leave excessive residues at harvest. However, Great Lakes Chemical assumes no responsibility, as to their accuracy nor for any loss due to excessive residues.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use.

Empty containers should be returned to Great Lakes Chemical Corp.

**TERR-O-GAS FUMIGATION DIRECTIONS
RECOMMENDED APPLICATION RATES**

Strawberries 400 to 500
lbs Acre

Expose to fumigation for 48 hrs. Aerate 2 weeks before setting transplants in treated areas.

Tomatoes 400 to 500
lbs Acre

Expose to fumigation for 48 hrs. Aerate 2 weeks before setting transplants in treated areas.

Apply with chisels spaced 12 inches apart to a depth of 8 inches. Seal fumigant with a drag or cultipacker immediately behind chisels. Cover with a gas-proof tarpaulin. Remove tarpaulin after recommended fumigation period and aerate soil for two weeks before planting. If odors persist disc or plow the soil to assist aeration.

Do not apply Terr-O-Gas 45 when there is little or no air movement.

Do not apply near buildings inhabited by humans or livestock or in areas adjacent to fields where valuable crops are growing.

NOTE: Vapors of Terr-O-Gas 45 are extremely irritating to the eyes. When applying or transferring to other containers, provide good ventilation. In the presence of discomforting concentrations of Terr-O-Gas 45 vapor wear a full face mask with an approved Black Canister for acid gas and organic vapors.

GREAT LAKES CHEMICAL CORPORATION
P. O. BOX 2200,
WEST LAFAYETTE, INDIANA 47906
19523 SOUTH SUSANA ROAD
COMPTON, CALIFORNIA 90221

MADE IN U.S.A.

ACCEPTED

TERR-O-GAS

PREPLANT SOIL FUMIGANT

ACTIVE INGREDIENTS:

Methyl Bromide
Chloropicrin

For control of Nematodes, Wireworms, Weed Grubs, Ants and Soil Borne Insects. (Example: ...)



DANGER
KEEP OUT OF REACH OF CHILDREN
POISON

DO NOT INHALE VAPORS • SEE ANTIDOTE & WARNING

Manufactured by
GREAT LAKES CHEMICAL CORPORATION
P. O. BOX 2200, WEST LAFAYETTE, INDIANA 47906

U.S. EPA Reg. No. 5000-21

TERR-O-GAS FUMIGATION DIRECTIONS
RECOMMENDED APPLICATION RATES

Berries 400 to 500
 lbs Acre
 before setting transplants in treated

Plants 400 to 500
 lbs Acre
 before setting transplants in treated

Apply with chisels spaced 12 inches apart to
 depth of 8 inches. Seal fumigant with a
 roller or cultipacker immediately behind
 chisel. Cover with a gas-proof tarpaulin.
 Remove tarpaulin after recommended fumi-
 gation period and aerate soil for two weeks
 before planting. If odors persist disc or plow
 to assist aeration.

Do not apply Terr-O-Gas 45 when there is
 any air movement.

Do not apply near buildings inhabited by
 humans or livestock or in areas adjacent to
 fields where valuable crops are growing.

Vapors of Terr-O-Gas 45 are extreme-
 ly irritating to the eyes. When apply-
 ing or transferring to other contain-
 ers, provide good ventilation. In the
 presence of discomforting concentra-
 tions of Terr-O-Gas 45 vapor wear
 a full face mask with an approved
 Black Canister for acid gas and
 organic vapors.

GREAT LAKES CHEMICAL CORPORATION
 P. O. BOX 2200,
 WEST LAFAYETTE, INDIANA 47906
 19523 SOUTH SUSANA ROAD
 COMPTON, CALIFORNIA 90221

ACCEPTED

June-17-1968

5785



TERR-O-GAS 45

PREPLANT SOIL FUMIGANT

ACTIVE INGREDIENTS:

Methyl Bromide 45%
 Chloropicrin 55%

By Wt.

Total 100%

For control of Nematodes, Wireworms, Weed Seeds, Fungus Diseases,
 Grubs, Ants and Soil Borne Insects. (Example: White Grubs)



DANGER

KEEP OUT OF REACH OF CHILDREN

POISON

DO NOT INHALE VAPORS • SEE ANTIDOTE & WARNINGS ON SIDE PANEL

Manufactured by

GREAT LAKES CHEMICAL CORPORATION

P. O. BOX 2200, WEST LAFAYETTE, INDIANA 47906

USDA Reg. No. 59624



POISON



DANGER

MAY BURN SKIN AND EYES
POISONOUS IF INHALED

Do Not Breathe Vapor
Do Not Get in Eyes, on Skin or Clothing

In case of contact, immediately flush skin or eyes
 with copious amounts of water for at least 15 minutes and get
 medical attention in case of contact. If liquid is spilled
 on clothing, remove clothing immediately, and
 do not rewear until completely clean and the liquid
 has been removed.

POISON

SEND FOR DOCTOR IN CASE OF ACCIDENT

ANTIDOTE: None known. In case of contact with
 liquid, wash with water. In case of contact with
 vapor, get to fresh air immediately. If
 breathing is difficult, give artificial respiration.
 If swallowed, give water and induce vomiting.

NOTICE

Following soil fumigation, undesirable concentra-
 tions of chloropicrin may drift to nearby areas. If
 this occurs, immediately cover treated area with a
 plastic tarpaulin. Leave tarpaulin on overnight and
 remove during the daytime. Recover areas if eye
 irritation indicates that undesirable concentrations
 are continuing. These concentrations are most likely
 to occur in the evening or at night when the air is
 most likely to be static in movement. Do not make
 application of Terr-O-Gas 45 when there is little or
 no air movement or there is an inversion condition in
 the atmosphere.

NOTE CAREFULLY: Following fumigation, the level
 of soluble salts and ammonia nitrogen may be
 raised. This is most likely to occur when heavy rates
 of fumigant and fertilizers are applied to soils that
 are either acid, wet, cold, or high in organic matter.
 Fertilizers containing ammonium salts should not be
 used. Apply only fertilizers containing nitrates until
 after the crop is well established and the soil tem-
 perature is above 65° F. in order to avoid ammonia
 injury, nitrate starvation, or both. After fumigation
 fertilize as indicated by soil test to protect against
 injury to plant roots. It is best that highly acid soils
 be limed before application to stimulate nitrification
 and reduce the possibility of ammonia toxicity.