



WARNING

KEEP AWAY from FEED or FOOD PRODUCTS

POISON

KEEP AWAY from FEED or FOOD PRODUCTS

**AUTOMATIC DO NOT DROP
DON'T**

DON'T

WARNING

Mr. Weston had no trouble at all in getting the men to do their work, and the result was that the steel minimum wage was established without difficulty, and the product of the miners' labour was increased.

FERR-O-GAS FUMIGATION DIRECTIONS

RECOMMENDED APPLICATION RATES

Strawberries 350 to 420
lbs Acre

Expose to fumigation for 48 hrs. Aerate 2 weeks before setting transplants in treated areas.

Tomatoes 350 to 420
bs/Acre

Expose to fumigation for 48 hrs. Aerate 2 weeks before setting transplants in treated areas

Apply, with chisels spaced 12 inches apart to a depth of 8 inches. Seal fumigant with a drag or cultipacker immediately behind chisels. Cover with a gas-proof tarpaulin. Remove tarpaulin after recommended fumigation period and aerate soil for two weeks before planting. If odors persist disc or plow the soil to assist aeration.


Do not apply Terr-O-Gas 57 when there is little or no air movement

Do not apply near buildings inhabited by humans or livestock or in areas adjacent to fields where valuable crops are growing.

NOTE: Vapors of Tarr-O-Gas® 57 are extremely irritating to the eyes. When applying or transferring to other containers, provide good ventilation. In the presence of disorienting concentrations of Tarr-O-Gas® vapor wear a full face mask with an approved black Canister filter and charcoal against vapors.

GREAT LAKES CHEMICAL CORPORATION

P. O. BOX 2700
WEST LAFAYETTE INDIANA 47906
INDIANA UNIVERSITY ROAD
MICHIGAN AVENUE




TERR-O-GA

PREPLANT SOIL FUL

ACTIVE INGREDIENTS:

Methyl Bromide.....
Chloropicrin

For control of Nematodes, Wireworms, Weed Seeds, Grubs, Ants and Soil Borne Insects (Example: White



ANSWER TO THE QUESTION OF THE DAY

—
—
—

1. *Leucosia* *leucostoma* *leucostoma* *leucostoma* *leucostoma*

GREAT LAKES CHEMICAL CORP
P. O. BOX 1100 - WELLS LA FAYETTE IN

**RR-GAS FUMIGATION DIRECTIONS
RECOMMENDED APPLICATION RATES**

berries 350 to 420
 lbs Acre
to fumigation for 48 hrs. Aerate 2
before setting transplants in treated

es 350 to 420
'bs/Acre
to fumigation for 48 hrs. Aerate 2
before setting transplants in treated

with chisels spaced 12 inches apart to
a depth of 8 inches. Seal fumigant with a
cultipacker immediately behind
the chisel plow. Cover with a gas-proof tarpaulin.
Remove tarpaulin after recommended fumi-
gation period and aerate soil for two weeks
before planting. If odors persist disc or plow
soil to assist aeration.

apply Terr-O-Gas 57 when there is no air movement

• apply near buildings inhabited by
• or livestock or in areas adjacent to
where valuable crops are growing.

Vapors of Terr-O-Gas 57 are extremely irritating to the eyes. When applying or transferring to other containers, provide good ventilation. In the presence of discomforting concentrations of Terr-O-Gas 57 vapor wear a full face mask with an approved Black Canister for acid gas and organic vapors.



TERR-O-GAS 57

PREPLANT SOIL FUMIGANT

ACTIVE INGREDIENTS:

By Wt

For control of Nematodes, Wireworms, Weed Seeds, Fungus Diseases, Grubs, Ants and Soil Borne Insects (Example: White Grubs)



KEEP OUT OF REACH OF CHILDREN

PUISSEN

Manufactured by

GREAT LAKES CHEMICAL CORPORATION

P. O. BOX 2200 WEST LAFAYETTE INDIANA 47906

REFERENCES



POISON

DANGER

MAY BURN SKIN AND EYES
POISONOUS IF INHALED

Do Not Breathe Vapor
Do Not Get in Eyes, on Skin or Clothing

10. The following table shows the number of hours worked by each of the 100 workers in the sample.

F-50N
SEND FOR DIRECTOR IN CASE OF ACCIDENT

NOTICE

Folk-wire soft fumigation, undesirable concentrations of chlorophacin may drift to nearby areas. If this occurs, immediately cover treated area with a plastic tarpaulin. Leave tarpaulin on overnight and remove during the daytime. Recover areas if eye irritation indicates that undesirable concentrations are continuing. These concentrations are most likely to occur in the evening or at night when the air is most likely to be static or movement. Do not make applications of TemOcar 57 when there is little or no air movement or there is an inversion condition in the atmosphere.

NOTE CAREFULLY: Following fumigation, the level of soluble salts and ammonia nitrogen may be raised. This is most likely to occur when heavy rates of fumigant and fertilizers are applied to soils that are either acid, wet, cold, or high in organic matter. Fertilizers containing ammonium salts should not be used. Apply only fertilizers containing nitrates until after the crop is well established and the soil temperature is above 65° F. in order to avoid ammonia injury, nitrate starvation, or both. After fumigation, apply an adequate amount of topsoil to protect against erosion. If the soil is dry, irrigate it frequently to prevent cracking and to encourage infiltration. If the soil is very sandy, fumigation