




Handwritten: 3-20-50
3-20-50
3-20-50

NOTICE

Many pesticidal chemicals are poisons and may leave a toxic residue on the plants to which they are applied. The U. S. Food and Drug Administration has established maximum amounts of such pesticidal chemicals that may remain on raw agricultural products at harvest, and it is the grower's responsibility to see that there is no residue on such crops at harvest in excess of these amounts. The "Fumigation Directions" are based on the best available information, and if followed carefully should not leave excessive residue at harvest. However, Great Lake Chemical assumes no responsibility for the residue on crops at harvest due to excessive use.

Seller warrants that the product contains the ingredients specified and is suitable for the purpose stated on the label when used in accordance with the directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product, or to any use not in strict accordance with the directions on the label, or to any use not in strict accordance with the label.

Empty containers should be returned to Great Lake Chemical Corporation, P. O. Box 100, Arkansas, via freight, collect.



WARNING

KEEP AWAY FROM FEED OR FOOD TRAYS

POISON

CAUTION—DO NOT DROP

DON'T

TERR-O-GAS[®] 75

PREPLANT SOIL FUMIGANT

ACTIVE INGREDIENTS:

	By Wt.
Methyl bromide	75%
Chloropicrin	25%
Total	100%

For control of nematodes, wireworms, weed seeds, fungus diseases, grubs, ants and soil borne insects. (Example: White grubs)




DANGER

KEEP OUT OF REACH OF CHILDREN

POISON

DO NOT INHALE VAPORS • SEE ANTIDOTE & WARNINGS ON SIDE PANEL



NET WEIGHT _____ LBS.

EPA Reg. No. 5785-40

Expose to fumigation for 48 hours. Aerate 2 weeks before setting transplants in treated areas.

Apply with chisels spaced 12 inches apart to a depth of 8 inches. Seal fumigant with a drag or cultipacker immediately behind chisel. Cover with a gas proof tarpaulin. Remove tarpaulin after recommended fumigation period and aerate soil for 2 weeks before planting. If odors persist, disc or plow the soil to assist aeration.

Do not apply Terr O Gas 75 when there is little or no air movement.

Do not apply near buildings inhabited by humans or livestock or in areas adjacent to them where valuable crops are cultivated.

NOTE: Vapors of Terr O Gas 75 are extremely irritating to the eyes. When applying in greenhouses or other containers, provide adequate ventilation. In the presence of high fumigant concentrations, use Terr O Gas 75 only in a well-ventilated tank with an exhaust system to remove fumigant vapors.

contact. If liquid is spilled, immediately remove, remove them immediately, and thoroughly. Do not reuse shoes or clothing that have been contaminated.

POISON

SEND FOR DOCTOR IN CASE OF POISONING. **ANTIDOTE:** Remove patient to fresh air and administer first aid as recommended. Hospitalize patient. Do not induce vomiting. Do not give anything by mouth.

NOTICE

Following soil fumigation, undesirable concentrations of chloropicrin may drift to nearby areas. If this occurs, immediately cover treated area with a plastic tarpaulin. Leave tarpaulin on overnight and remove during the daytime. Recover the area if eye irritation indicates that the undesirable concentration is continuing. These concentrations are most likely to occur in the evening or at night when the air is most likely to be static in movement. Do not make application of Terr O Gas 75 when there is little or no movement or there is an inversion condition in the atmosphere.

NOTE CAREFULLY: Following fumigation, the level of soluble salts and ammonia nitrogen may be raised. This is most likely to occur when heavy rates of fumigant and fertilizers are applied to soils that are either acid, wet, cold, or high in organic matter. Fertilizers containing ammonium salts should not be used. Apply only fertilizers containing nitrates until after the crop is well established and the soil temperature is above 65° F. in order to avoid ammonia injury, nitrate starvation, or both. After fumigation, fertilize as indicated by soil test to protect against injury to plant roots. It is best that highly acid soils be limed before application to stimulate nitrification and reduce the possibility of ammonia toxicity.