



TERR-O-CIDE 93.5

For preplant soil fumigation to control insects, nematodes, weeds and fungi

Active Ingredients 100%

Chloropicrin	0.3	5
Ethylene dibromide	6	5
Total	100	0

DANGER

KEEP OUT OF REACH OF CHILDREN

PAPERS


**POISONOUS IF INHALED, SWALLOWED
OR CONTACTED WITH SKIN**

REFERENCES

6. $\{x_1, x_2, \dots, x_n\} \subseteq \{1, 2, \dots, n\}$

ANNUAL ANTHROPOLOGICAL

NOTE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.



GENERAL INFORMATION

Terr-O-Cide 93.5% Chloropicrin is used for control of the grass, weeds and fungi to be planted to millet, rye, sweet corn, sugar cane, eggplant, etc., prior to planting in nursery, to be planted to strawberries and potato beds, to be planted to asparagus, artichokes, etc., to pasture or to berry bushes, etc., or to control certain nematodes, fungi, viruses, etc. After the fumigant Terr-O-Cide 93.5% vaporizes from the soil surface, it volatizes and diffuses through the soil controlling most of the fungi causing damping off, root rot, wilt, etc., bacteria, soil insects, nematodes, weeds and weed seeds. (Especially recommended when completely rotten root galls are present.)

DIRECTIONS FOR USE

All types of equipment, from hand injectors to tractor drawn machines, are used for the application of Terr-O-Cide 93.5% vapor. Depending upon the size of the area treated, effectiveness of treatment increases proportionately. Treat at temperatures above 60°F. For best kill of weed seed, treat and wait at least four days prior to fumigation.

DOSAGE Normal dosage is 2-3 cubic feet per 10' centers in light soil; 8-9 cubic feet for heavier soils. When using a hand-type applicator, inject 6-10 pint per 100 linear feet at 12" spacing. Inject 4 to 6" deep or at half-bench digging. Use deeper injections in light soils or at higher temperatures. Consult your State Agricultural Extension Service for local conditions.

For bulk soil treatment use 1-2-3-4 lb per cubic yard depending on texture.
GAS RETENTION For small areas take over injection points, immediately, and seal with water seal. For best results, use a polyethylene tape cover. For larger areas, seal plowed soil immediately by leveling and pulling a tarp or plastic sheet over it. Half acre inch of water added to a quarter of an acre improves soil infiltration. Laying of polyethylene tarp is used at the edges of the treated areas for moisture control.

EXPOSURE PERIOD Expose 11-18 days according to 1% vapor level in the bulk soil.

AERATION Do not plant until the vapor content has dropped below 1% vapor level. Higher vapor levels over longer periods of time will require a longer time for fumigation.

PRECAUTIONS

Terr-O-Cide 93.5% Chloropicrin is extremely dangerous to humans and animals and bulky from every aspect. It is a volatile liquid which can easily escape from some metal drums, but tends to stay in plastic containers. It is easily absorbed from the skin. If fumigant vapors come in contact with the eyes, wash immediately with cool, but not cold, water. If fumigant vapors come in contact with the skin, wash immediately with soap and water. If fumigant vapors come in contact with the lungs, remove to fresh air. If fumigant vapors come in contact with the mouth, drink water.

DIRECTIONS FOR EMPTYING CYLINDER

Do not cut cylinder open. Do not pour contents of cylinder onto ground. Do not content around pets or people.

Remove the large valve from the cylinder. Turn cylinder upside down and pour the liquid into a tank truck, etc. Turn cylinder right side up and pour the remaining liquid into a tank truck, etc. Turn cylinder upside down again and pour the remaining liquid into a tank truck, etc.

Do not cut cylinder open. Do not pour contents of cylinder onto ground. Do not content around pets or people.

STORAGE AND RETURN OF CYLINDER

ERR-O-CIDE 93.5 SOIL FUMIGANT

For preplant soil fumigation to control insects, nematodes, weeds and fungi

Active Ingredients 100%

Chloropicrin	93.5
Ethylene dibromide	6.5
Total	100.0

DANGER

KEEP OUT OF REACH OF CHILDREN
POISON

POISONOUS IF INHALED, SWALLOWED
OR CONTACTED WITH SKIN

EPA Reg. No. 5785-44

FENT

NOTE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.