

Sept - 25-1971

5777-164

DIRECTIONS FOR USE:

CROP	INSECTS	RATE	REMARKS
Tobacco	Homworms Flea Beetles	2 Qts. per acre.	Apply in 20-25 Gallons of water per acre. Do not use within 5 days of priming or 15 days of cutting.

Use of this material in a manner or at a time other than in accordance with the directions may produce plant injury, excessive residues or other undesirable results.

Rinse equipment and containers and dispose of wastes by leaving in non-crop lands away from water supplies. Containers should be disposed by punching holes in them and burying with wastes.

NOTICE:

See page 10 for complete directions. Do not use product more than indicated on the label. Do not use on food crops unless specifically labeled for such use. Do not use on food crops contrary to label directions.

SUPER FLASH TOBACCO INSECTICIDE

ACTIVE INGREDIENTS:

O,O-Dimethyl S-4-oxo-1,2,3-benzotriazin-3(4H)-ylmethyl phosphorodithioate*	12.37%
Xylene	83.63%
INERT INGREDIENTS	4.00%
TOTAL	100.00%

*U. S. Patent No. 2,753,115

(See side panels for warnings and antidote)

Manufactured By
CAROLINA CHEMICAL CORP.
Wilson, N. C.

SUPER FLASH **BACCO INSECTICIDE**

INGREDIENTS:

ethyl S-4-oxo-1,2,3-benzotriazin-3 (4H)	
ethyl phosphorodithioate*	12.37*
.....	83.63%
INGREDIENTS	4.00%
TOTAL	100.00%

Patent No. 2,758,115

(See side panels for warnings and antidote)

Manufactured By
CAROLINA CHEMICAL CORP.
Wilson, N. C.

USDA P. 22-311-103