

# FOGICIDE-X

SPACE SPRAY INSECTICIDE

A high level, synergized pyrethrin type insecticide concentrate for space application by fogging and spraying machines and similar equipment. FOGICIDE XL can be used around food, food products, utensils and equipment provided that no exposed food is present and provided that food utensils and surfaces that will be in contact with food are covered and provided that fogging and spraying is not performed in food processing plants while plant is in operation.

Properly applied, FOGICIDE XL kills the accessible exposed stages of flies, ants, mosquitos, boxelder bugs, silverfish flying moths, gnats, crickets, roaches, spiders, scorpions, wasps, cornets, clover mites, cheese mites, cheese skipper, and most common stored-grain pests, including granary weevil, rice weevil, grain beetle, spider beetle, angoumois grain moth, Mediterranean flour moth and grain mites.

Active INGREDIENTS

- \*Piperonyl Butoxide, Technical Pyrethrins
- Petroleum Oil
- \*Equivalent to 1/6% of (butylcarbityl) (6-propylpiperonyl) ether and to 0.4% related compounds.

USDA Registration No. 5870-10



## CAUTION TREAD BEFORE USING

### HARMFUL IF SWALLOWED KEEP OUT OF REACH OF CHILDREN

Do not remain in treated areas and ventilate the area after treatment is completed. FOGICIDE XL is an oil base material and should not be sprayed, stored or otherwise exposed to an open flame.

In order to avoid any possible explosive hazard, not more than 1 gallon should be sprayed per 40,000 cu. ft. of enclosed space.

Remove or cover all exposed foods or foodstuffs. Cover all utensils and surfaces that will be in contact with food. Do not spray or fog in food processing plant while plant is in operation.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in safe place.

## FOGICIDE-XL

SPACE SPRAY INSECTICIDE

OGICIDE XL kills the accessible exposed mosquitos, boxelder bugs, silverfish flying, roaches spiders, scorpions, wasps, cornets, mites, cheese skipper, and most common acluding granary weevil, rice weevil, grain e, angournois grain moth, Mediterranean mites.

Active INGREDIENTS

\*Piperonyl Butoxide, Technical Pyrethrins


Petroleum Oil

\*Equivalent to 1/6% of (butylcarbityl) (6-propylpiperonyl) ether and to 0.4% related compounds.

Dept 30 1971

100.0% 2.0% 0.20% 97.80% For efficient results, it is important that the insecticides be fogged or sprayed uniformly per unit of space. For crawling insects, use 2 ounces/1,000 cu. ft. and 1 ounce/1,000 cu. ft. from flying insects. If possible, keep area closed for 15-20 minutes after application.

USDA Registration No. 5870-10





### CAUTION: IREAD BEFORE USING

#### HARMFUL IF SWALLOWED KEEP OUT OF REACH OF CHILDREN

Do not remain in treated areas and ventilate the area after treatment is completed.

FOGICIDE XL is an oil base material and should not be sprayed, stored or otherwise exposed to an open flame.

In order to avoid any possible explosive hazard, not more than 1 gallon should be sprayed per 40,000 cu. ft. of enclosed space.

Remove or cover all exposed foods or foodstuffs. Cover all utensils and surfaces that will be in contact with food. Do not spray or fog in food processing plant while plant is in operation.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in safe place.

Active INGREDIENTS

\*Piperonyl Butoxide, Technical

**Pyrethrins** 

osed

lying

nets,

nome

grain.

nean

'RAY INSECTICIDE

Petroleum Oil

\*Equivalent to 1/6% of (butylcarbityl) (6-propylpiperonyl) ether and to 0.4%

related compounds.

gistration No. 5870-10

100.0%

2.0%

0.20% 97.80%

For efficient results, it is important that the insecticides by fogged or sprayed uniformly per unit of space. For crawling insects, use 2 ounces 1,000 cu. ft. and 1 ounce 1,000 cu. ft. f  $\sim$ flying insects. If possible, keep area closed for 15-20 minu is

- ¿\*.

after application.