

COLOROX SERVICE BULLETIN

COLOROX (5.25% sodium hypochlorite, EPA REG. 5813-1)

VIRUCIDAL EFFICACY

Clorox, when used as directed below, is effective against the following viruses:

Adenovirus Type 2

Hepatitis A

Human Immunodeficiency Virus

Respiratory syncytial virus

Herpes simplex virus 2

Rubella virus

*Feline parvovirus**

Rotavirus

Cytomegalovirus

Influenza A2

Varicella zoster virus

Rhinovirus Type 17

*Canine parvovirus**

Directions for use:

Hard Non-porous Surfaces:

To disinfect hard non-porous surfaces, clean surface by removing gross filth (loose dirt, debris, food materials, etc.). Prepare a solution of 3/4 cup (6 oz.) Clorox per gallon of water (~2400ppm available chlorine). Thoroughly wet surface with the solution and allow it to remain in contact with the surface for 5 minutes. Rinse with clean water and dry.

* For Canine and Feline parvovirus use the same instructions as above but keep the solution in contact with the surface for 10 minutes.

Toilet Bowls:

Flush toilet to remove gross filth. Add 1 cup Clorox to the bowl and brush surfaces thoroughly. Let stand 5 minutes before flushing again.

Use Sites

Clorox can be used on hard non-porous surfaces in commercial, institutional, hospital and household premises (including kitchens, bathrooms, nurseries, sick rooms, laundry rooms, garbage containers and diaper pails), eating establishments, pet kennels and veterinary premises.

The Clorox Company
1221 Broadway
Oakland, CA 94615

NOV 22 1994

BEST AVAILABLE COPY

5813-1

2 7 3

CLOROX SERVICE BULLETIN

CLOROX (5.25% sodium hypochlorite, EPA REG. 5813-1)

BACTERICIDAL EFFICACY

Clorox, when used as directed below, is effective against the following bacteria:

Gram positive bacteria including:

Staphylococcus aureus

Streptococcus pyogenes

Gram negative bacteria including:

Salmonella choleraesuis

Pseudomonas aeruginosa

Escherichia coli serotype 0157:H7

Legionella pneumophila

Klebsiella pneumoniae

Directions for use:

Hard Non-porous Surfaces:

To disinfect hard non-porous surfaces, clean surface by removing gross filth (loose dirt, debris, food materials, etc.). Prepare a solution of 3/4 cup (6 oz.) Clorox per gallon of water (2400ppm available chlorine). Thoroughly wet surface with the solution and allow it to remain on the surface for 5 minutes. Rinse with clean water and dry.

Toilet Bowls:

Flush toilet to remove gross filth. Add 1 cup Clorox to the bowl and brush surfaces thoroughly. Let stand 5 minutes before flushing again.

Use Sites

Clorox can be used on hard non-porous surfaces in commercial, institutional, hospital and household premises (including kitchens, bathrooms, nurseries, sick rooms, laundry rooms, garbage containers and diaper pails), eating establishments, pet kennels and veterinary premises.

The Clorox Company
1221 Broadway
Oakland, CA 94615

BEST AVAILABLE COPY

CLOROX SERVICE BULLETIN

CLOROX (5.25% sodium hypochlorite, EPA REG. 5813-1)

TUBERCULOCIDAL EFFICACY

Clorox, when used as directed below, is effective against *Mycobacterium bovis*

Directions for use:

Hard Non-porous Surfaces:

To disinfect hard non-porous surfaces, clean surface by removing gross filth (loose dirt, debris, food materials, etc.). Prepare a dilution of 1 part Clorox to 10 parts water (13 oz/gallon.) (~5000ppm available chlorine). Thoroughly wet surface with the solution and allow it to remain in contact with the surface for 5 minutes. Rinse with clean water and dry.

Use Sites

Clorox can be used on hard non-porous surfaces in commercial, institutional, hospital and household premises (including kitchens, bathrooms, nurseries, sick rooms, laundry rooms, garbage containers and diaper pails), eating establishments, pet kennels and veterinary premises.