

green meadow

WEED KILLER

AND

LAWN FOOD

CAUTION

Read and Observe Precautions
on Back Panel for Additional Cautions
DANGER TO HUMANS

COVERS 5000 SQ. FT.


NON-BURNING
WHEN USED AS DIRECTED

Facit Label


*MM
MM
MM*

*LAN
LAW
LAW*

NET WEIGHT 20 POUNDS



**WEED KILLER
AND
LAWN FOOD
10-6-4
DIRECTIONS**



APPLYING

Product should be applied with a good spreader at rate of 4 lbs. per 1000 square feet. Avoid overlap of spreading by using some type of marker to indicate covered area.

TIME OF APPLICATION

Most broad leaf weeds will be controlled by applying when ground is moist and weeds are growing rapidly. For best results, apply several days after mowing. Then do not cut grass for several more days.

Controls these and many other broadleaf weeds:
Purslane - Dandelions Buckhorn - Plantain

FOLLOW-THRU

Sometimes weed infestation will result a complete kill. A follow-up application is desirable if desired results are not obtained in 6 to 8 weeks after first application and will also control new weed growth.

CAUTION

- 1 - Apply on calm days.
- 2 - Thoroughly clean all equipment used.
- 3 - Do not use on bent or clover.
- 4 - Do not store near other fertilizer, seed, insecticides or fungicides.
- 5 - Heavy applications than recommended may be harmful.

Avoid taking dust and dirt to other plants and crops. 6 - Never apply on new seed beds or to newly planted grasses until after several months of growth. Avoid contact with skin, eyes or clothing. Contamination of water for domestic use.

Do not reuse container. Destroy when empty.

SPREADER SETTINGS

Cyclone	4 1/4	Lawn Beauty	6
Scotts	6 1/2	Cross Country (Seas) ..	7

Manufactured by

STADLER FERTILIZER
GRAND RAPIDS, OHIO 43522

Handwritten note: 6-10-72