


NET WEIGHT 20 POUNDS



*SPARK-GREEN*



**WEED KILLER  
AND  
LAWN FOOD  
10-6-4**

**MINIMUM GUARANTEED ANALYSIS**

Total Nitrogen . . . . .	10.00%
9.42% Ammoniacal Nitrogen	
.58% Nitrate Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) . . . . .	6.00%
6.00% Total Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	
Soluble Potash (K <sub>2</sub> O) . . . . .	4.00%
Potential Acidity Equivalent to 925 lbs. of Calcium Carbonate per ton.	

ACTIVE INGREDIENTS  
 1.00% Active Salt of 2,4-Dichlorophenoxyacetic Acid\*  
 INERT INGREDIENTS  
 99.00%  
 \*Equivalent to 2,4-Dichlorophenoxyacetic Acid 63%  
**TOTAL 100.00%**

# DIRECTIONS

## APPLYING

Product should be applied with a good spreader at rate of 4 lbs. per 1000 square feet. Avoid overlap of spreading by using some type of marker to indicate covered area.

## TIME OF APPLICATION

Most broad leaf weeds will be controlled by applying when ground is moist and weeds are growing rapidly. For best results, apply several days after mowing. Then do not cut grass for several more days.

Controls these and many other broadleaf weeds:  
Purslane • Dandelions  
Buckhorn • Plantain

## FOLLOW-THRU

Sometimes weed infestation will resist a complete kill. A follow-up application is desirable if desired results are not obtained in 6 to 8 weeks after first application and will also control new weed growth.

- 1 - Apply on calm days.
- 2 - Thoroughly clean all equipment used.
- 3 - Do not use on bent or clover.
- 4 - Do not store near other fertilizer, seed, insecticides or fungicides.
- 5 - Heavier applications than recommended may be harmful.

Avoid  
Inhaling dust,  
Wind drift to other plants  
and crops.

6 - Never apply in new seed beds or to newly planted grasses until after several months of growth.

Avoid:  
Contact with skin, eyes, or  
clothing. Contamination of  
water for domestic use.

## SPREADER SETTINGS

Stadler's Triple XXX.....8	Lawn Beauty.....6
Scotts.....6½	Cross Country (Sears)....7

Manufactured by

**THE STADLER FERTILIZER CO.**

CLEVELAND, OHIO 44109

## CAUTION

Keep Out of Reach of Children  
(See Back Panel for Additional Cautions)

Total Nitrogen - 10.00% (9.42% Ammoniacal Nitrogen, .58% N<sub>2</sub> Nitrogen)  
Available Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>) - 6.00% (6.00% Total Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>)  
Soluble Potash (K<sub>2</sub>O) - 4.00% (Potential Acidity Equivalent to 928 lbs. of  
Calcium Carbonate per ton)

ACTIVE INGREDIENTS: (Dimethylamine Salt of 2,4-Dichlorophenoxyacetic  
Acid) - 1.00%

INERT INGREDIENTS: 29.00% (Equivalent to 2,4-Dichlorophenoxyacetic Acid  
Salt) TOTAL - 100.00%

U.S.D.A. REG. NO.


# COVERS 5000 SQ. FT.

## NON-BURNING (WHEN USED AS DIRECTED)

5815-1  
ACCEPTED  
May 27 1967  
GENERAL INSECTICIDE  
REGISTERED  
SUBJECT  
COMMENTS



NET WEIGHT 20 POUNDS



PROFESSIONAL  
**WEED KILLER and LAWN FOOD**  
**10-6-4**

**MINIMUM GUARANTEED ANALYSIS**

Total Nitrogen . . . . . 10.00%  
9.42% Ammoniacal Nitrogen  
.58% Nitrate Nitrogen  
Available Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>) . . . . . 6.00%  
6.00% Total Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>)  
Soluble Potash (K<sub>2</sub>O) . . . . . 4.00%  
Potential Acidity Equivalent to 925 lbs. of Calcium  
Carbonate per ton.

ACTIVE INGREDIENTS:  
Dimethyl Amine Salt of 2,4-Dichlorophenoxyacetic Acid\* 1.00%  
INERT INGREDIENTS: 99.00%  
\*Equivalent to 2,4-Dichlorophenoxyacetic Acid .83% of 1%  
TOTAL 100.00%

ACCEPTED  
MAY 27 1967  
5415-7

**CAUTION**

Keep Out of Reach of Children  
(See Back Panel for Additional Cautions)

Total Nitrogen - 10.00% (9.42% Ammoniacal Nitrogen, .58% Nitrate Nitrogen)  
Available Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>) - 6.00% (6.00% Total Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>))  
Soluble Potash (K<sub>2</sub>O) - 4.00% (Potential Acidity Equivalent to 925 lbs. of Calcium Carbonate per ton)  
ACTIVE INGREDIENTS: (Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid\*) - 1.00%  
INERT INGREDIENTS: 99.00% (\*Equivalent to 2,4-Dichlorophenoxyacetic Acid .83% of 1%) TOTAL - 100.00%

U.S.D.A. REG. NO. [REDACTED]

**COVERS 5000 SQ. FT.**



NET WEIGHT 2

*SPARK-L*



**WEED K**

AND

**LAWN F**

**10-6**

**MINIMUM GUARAN**

Total Nitrogen . . . . .

9.42% Ammoniacal Nitrogen

.58% Nitrate Nitrogen

Available Phosphoric Acid (P)

6.00% Total Phosphoric A

Soluble Potash  $K_2O$  . . . . .

Potential Acidity Equivalent

Carbonate per ton.

AMERICAN FERTILIZER COMPANY  
24-0  
E. B. BROWN



NET WEIGHT 20 POUNDS



# TRIPLE GREEN

PROFESSIONAL  
**WEED KILLER and LAWN FOOD**  
**10-6-4**

### MINIMUM GUARANTEED ANALYSIS

Total Nitrogen . . . . .	10.00%
9.42% Ammoniacal Nitrogen	
.58% Nitrate Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) . . . . .	6.00%
6.00% Total Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> )	
Soluble Potash (K <sub>2</sub> O) . . . . .	4.00%
Potential Acidity Equivalent to 925 lbs. of Calcium Carbonate per ton.	

ACTIVE INGREDIENTS  
Dimethyl Amine Salt of 2,4-Dichlorophenoxyacetic Acid 99.00%  
INERT INGREDIENTS  
\*Equivalent to 2,4-Dichlorophenoxyacetic Acid 1.00%  
TOTAL 100.00%

### CAUTION

Keep Out of Reach of Children  
see Back Panel for Additional Information

Total Nitrogen 10.00% Ammoniacal Nitrogen 9.42% Nitrate Nitrogen .58%  
Available Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>) 6.00% Total Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>) 6.00%  
Soluble Potash (K<sub>2</sub>O) 4.00% Potential Acidity Equivalent to 925 lb. Calcium Carbonate per ton.  
ACTIVE INGREDIENTS: Dimethylamine Salt of 2,4-Dichlorophenoxyacetic Acid 99.00%  
INERT INGREDIENTS: 1.00%  
TOTAL 100.00%

U.S.D.A. REG. NO. [REDACTED]

## COVERS 5000 SQ. FT.

# DIRECTIONS

## APPLYING

Apply this product with a broadcast spreader. Apply to the lawn in a regular pattern, overlapping each row.

## TIME OF APPLICATION

Most lawn weeds will be controlled by one application. A second application at 6 to 8 weeks after the first application will control weeds that have not been controlled by the first application. For several more days.

Control these and many other broadleaf weeds:  
Purslane • Garden Spurge • Buckhorn • Knotweed

## FOLLOW-THRU

Sometimes weed infestation will resist a complete kill. A follow-up application is desired if desired results are not obtained in 6 to 8 weeks after first application and will also control new weed growth.

## CAUTION

- 1—Apply on calm days
- 2—Thoroughly clean all equipment used.
- 3—Do not use on bent or crever
- 4—Do not store near other fertilizer, seed, insecticides or fungicides.
- 5—Heavier applications than recommended may be harmful.

Apply to new seed beds or to newly planted grasses until after several months of growth.

Apply to new seed beds or to newly planted grasses until after several months of growth.

Apply to new seed beds or to newly planted grasses until after several months of growth.

Apply to new seed beds or to newly planted grasses until after several months of growth.

Apply to new seed beds or to newly planted grasses until after several months of growth.

## SPREADER SETTINGS

Stadler's Triple XXX.....8	Lawn Beauty.....6
Scotts.....6½	Cross Country (Sears)....7

A PRODUCT OF SPEE-D-GRO COMPANY, CLEVELAND 9, OHIO