



STEIN SUPREME

Weed & feed

HERBICIDE
ACTIVE INGREDIENTS
 *Dimethylamine salt of 2,4-dichlorophenoxyacetic acid 1.41%
 **Isocetyl ester of 5-ters. 2-12-15 trichloro-oxo) propionic acid 0.20%
INERT INGREDIENTS 98.39%
TOTAL 100.00%

*Equivalent to 2,4-dichlorophenoxyacetic acid 1.83%
 **Equivalent to (2-(2,4,5-trichlorophenoxy) propionic acid 0.24%

EPA Reg. No. 5815-31

22-6-6

ACCEPTED
 12/6/75
 U.S. DEPARTMENT OF AGRICULTURE
 OFFICE OF PESTICIDE REGISTRATION
 WASHINGTON, D.C. 20250

CONTROLS DANDELIONS,
 CHICKWEED, CLOVER

FERTILIZES THE LAWN

COVERS 5000 sq. ft.

Weed & feed

HERBICIDE

ACTIVE INGREDIENTS

**Methoxychlor	1.3%
dichlorophenoxyacetic acid	0.83%
**Isopropyl ester of Silver 2-(2,4,5-trichlorophenoxy) propionic acid	0.14%

INERT INGREDIENTS 98.80%

TOTAL 100.00%

*Equivalent to 2,4-dichlorophenoxyacetic acid 0.83%

**Equivalent to 2-(2,4,5-trichlorophenoxy) propionic acid 0.14%

EPA Reg. No. 5815-31

22-6-6

ACCEPTED

12/6/75

UNDER THE FEDERAL INSECTICIDE ACT

5815-31

**CONTROLS DANDELIONS,
CHICKWEED, CLOVER**

FERTILIZES THE LAWN

COVERS 5000 sq. ft.



CAUTION

See label for directions and first aid instructions.

20 LB. NET WEIGHT



STEIN SUPREME weed & feed

*GUARANTEED FERTILIZER ANALYSIS	
Total Nitrogen (N)	22.00%
Available Phosphoric Acid (P ₂ O ₅)	6.00%
Soluble Potash (K ₂ O)	6.00%

*In Milwaukie area only.

SUPREME WEED & FEED HAS BEEN SCIENTIFICALLY FORMULATED TO ASSURE A SUPERIOR LUSH-GREEN-WEED-FREE LAWN.

SUPREME WEED & FEED CONTAINS THE BASIC PLANT FOOD ELEMENTS FOR PLANT GROWTH. THE NUTRIENTS ARE IN PROPER PROPORTION TO MEET THE TURF NEEDS OF LAWNS IN THE MILWAUKEE AREA.

SUPREME WEED & FEED HAS BUILT IN WEED KILLERS CONSISTING OF 2, 4-D, AND SILVEX. THESE WORK TOGETHER IN CONTROLLING UNDESIRABLE BROADLEAF WEEDS.

DIRECTIONS

APPLICATION SHOULD BE MADE WHEN WEEDS ARE YOUNG AND ACTIVELY GROWING.

WEED KILLERS WORK THRU THE FOLIAGE OF THE WEED. THEREFORE FOR GUARANTEED RESULTS THIS PRODUCT **MUST BE APPLIED TO WET GRASS**. EARLY MORNING OR LATE EVENING APPLICATIONS WHEN TURF IS WET WITH DEW WOULD BE IDEAL. IF APPLIED DURING THE DAY, SPRINKLE LIGHTLY BEFORE APPLICATION.

DO NOT MOW OR WATER THE LAWN AT LEAST 24 HOURS AFTER APPLICATION.

Set the spreader as shown below. Contents of this bag covers 5,000 sq. ft.

See cautions below.

SUGGESTED SPREADER SETTINGS

4# PER 1,000 SETTING

SPREADER	SPREADER
AGRICO AGRI-SPRED	GREENFIELD, CENTRAL
JACKSON, SEARS	LAWN BEAUTY, M. T. D.
AGRICO JET SPRED, CYCLONE ..	SCOTTS

WALKING SPEED, EVENNESS OF LAWN AND CONDITION OF SPREADER MAY REQUIRE SLIGHTLY DIFFERENT SETTINGS FOR DESIRED COVERAGE.

CAUTION

Keep out of reach of children.

Avoid contact with skin, eyes and clothes.

Harmful if swallowed.

Avoid breathing dust.

Do not reuse container. Destroy when empty. Do not burn.

Do not store near other fertilizers, seeds, insecticides or fungicides.

Do Not use on Centipede, St. Augustine, Dichondra, or Carpet grasses. Temporary injury may be experienced on Bentgrass.

Keep application at least 2 feet away from shrubs, vegetable gardens, and sensitive trees (such as Willow, Locust, and Dogwood). If new growth on trees or shrubs starts to curl, water heavily.

If grass clippings from the first mowing after use of this product are removed, do not feed to livestock or use to mulch ornamental shrubs.

Do not use on newly seeded lawns.

Soluble Polymers (S.P.) 6.00%
* In 100% wet weight

SUPREME WEED & FEED HAS BEEN SCIENTIFICALLY FORMULATED TO ASSURE A SUPERIOR LUSH-GREEN-WEED-FREE LAWN.

SUPREME WEED & FEED CONTAINS THE BASIC PLANT FOOD ELEMENTS FOR PLANT GROWTH. THE NUTRIENTS ARE IN PROPER PROPORTION TO MEET THE TURF NEEDS OF LAWNS IN THE MILWAUKEE AREA.

SUPREME WEED & FEED HAS BUILT IN WEED KILLERS CONSISTING OF 2, 4-D, AND SILVEX. THESE WORK TOGETHER IN CONTROLLING UNDESIRABLE BROADLEAF WEEDS.

DIRECTIONS

APPLICATION SHOULD BE MADE WHEN WEEDS ARE YOUNG AND ACTIVELY GROWING.

WEED KILLERS WORK THRU THE FOLIAGE OF THE WEED. THEREFORE FOR GUARANTEED RESULTS THIS PRODUCT **MUST BE APPLIED TO WET GRASS.** EARLY MORNING OR LATE EVENING APPLICATIONS WHEN TURF IS WET WITH DEW WOULD BE IDEAL. IF APPLIED DURING THE DAY, SPRINKLE LIGHTLY BEFORE APPLICATION.

DO NOT MOW OR WATER THE LAWN AT LEAST 24 HOURS AFTER APPLICATION.
Set the spreader as shown below. Contents of this bag covers 5,000 sq. ft.
See cautions below.

SUGGESTED SPREADER SETTINGS

4# PER 1,000 SETTING

SPREADER		SPREADER	
AGRICO AGRI SPRED	5	GREENFIELD, CENTRAL	7
JACKSON, SEARS	5	LAWN BEAUTY, M.T.D.	5½
AGRICO JET SPRED, CYCLONE ..	4½	SCOTTS	5

WALKING SPEED, EVENNESS OF LAWN AND CONDITION OF SPREADER MAY REQUIRE SLIGHTLY DIFFERENT SETTINGS FOR DESIRED COVERAGE.

CAUTION

- Keep out of reach of children.*
- Avoid contact with skin, eyes and clothes.*
- Harmful if swallowed.*
- Avoid breathing dust.*
- Do not reuse container. Destroy when empty. Do not burn.
- Do not store near other fertilizers, seeds, insecticides or fungicides.
- Do not use on Centipede, St. Augustine, Dichondra, or Carpet grasses. Temporary injury may be experienced on Bentgrass.
- Keep application at least 2 feet away from shrubs, vegetable gardens, and sensitive trees (such as Willow, Locust, and Dogwood). If new growth on trees or shrubs starts to curl, water heavily.
- If grass clippings from the first mowing after use of this product are removed, do not feed to livestock or use to mulch ornamental shrubs.
- Do not use on newly seeded lawns.
- Vapors from this product may injure susceptible plants in the immediate vicinity.
- THIS PRODUCT IS TOXIC TO FISH. KEEP OUT OF LAKES, STREAMS OR PONDS. DO NOT APPLY WHERE RUNOFF IS LIKELY TO OCCUR. DO NOT CONTAMINATE WATER BY CLEANING OF EQUIPMENT, OR DISPOSAL OF WASTES. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.**

MANUFACTURED FOR
STEIN GARDEN CENTERS
MILWAUKEE, WISCONSIN