

JUN 10 1961

Mr. Dan Paradiso, Jr
The Andersons Lawn Fertilizer Division
d/d/a tree flow fertilizer
P.O. box 119
Maunee, WI 53537

Dear Mr. Paradiso:

Subject: Triple X Garden Weed Preventer
EPA Registration No. 5815-35
Application to Amend Registration, To Conform to
EPA Standard, June, 1968

The subject labeling submitted in connection with
registration under the Federal Insecticide, Fungicide and
Rodenticide Act (FIFRA) as amended is acceptable provided
that you:

1. Add the following statements under the heading:
"Directions For Use":

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE
CLOTHING:

Wear long-sleeved shirt and long-legged pants
and chemical resistant gloves.

IMPORTANT: If the pesticide comes in contact with
skin wash off with soap and water. Always wash
hands and face, and arms with soap and water before
smoking, eating, drinking, or toileting.

AFTER USE: Before removing gloves, wash them with
soap and water. Take off all work clothes and
shoes. Shower using soap and water. Do not reuse
contaminated clothing. Clothing worn during use
must be laundered separately from household articles.

BEST AVAILABLE COPY

2. Submit five (5) printed copies of the final printed label before releasing the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA, Section 6(e). Your release for shipment of the product constitutes acceptance of this condition.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

Joanne L. Miller
Product Manager (23)
Fungicide-Herbicide Branch
Registration Division (R-7505C)

Enclosure

E.Wilson:Diskette #0007b: 06-05-91

BEST AVAILABLE COPY

(front panel)

This bag is a 2-color
pinch-bottom bag with
black type printed on
a light blue background

TRIPLE X GARDEN WEED PREVENTER

COVERS 2,500 SQ. FT.

ACTIVE INGREDIENT

DPCA (dimethyl tetrachloroterephthalate). 5.00%

INERT INGREDIENTS 95.00%

TOTAL 100.00%

KEEP OUT OF REACH OF CHILDREN ~ 12 pt.

CAUTION ~ 18 pt.

If in Eyes: Flush with plenty of water.
Call a physician if irritation persists.

If on Skin: Wash with plenty of soap and water.
Call a physician if irritation persists.

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Reg. No. 5815-35

EPA Est. 9198-OH-1

10 POUNDS NET WT.

**ACCEPTED
with COMMENTS
in EPA Letter Dated:**

FOR 11-19-71

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, this pesticide
registered under EPA Reg. No.**

5815-35

BEST AVAILABLE COPY

(back panel)

TRIPLE X GARDEN WEED PREVENTER

KEEP OUT OF REACH OF CHILDREN

PRECAUTIONARY STATEMENTS

CAUTION

Hazards to Humans and Domestic Animals

Harmful if swallowed. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling. Avoid breathing dust. Do not contaminate food and feedstuffs. Do not store near seed, fertilizer or other herbicides.

ENVIRONMENTAL HAZARDS

Collect and incorporate granules spilled on the soil surface. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

~~Read the entire label carefully before using this product.~~

It is in violation of Federal Law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergence herbicide for the control of many broadleafed weeds and most annual grasses ~~on lawns and around~~ for use around vegetables, strawberries, home flower gardens and nursery stock.

Any application should be made prior to weed seed germination Recommended Rate: Apply at a rate of 4 pounds per 1,000 square feet or 1 pound per 250 square feet. Weed control up to three months can be expected following proper application. Application should be made to a uniform weed free soil within two or three days following cultivation to prevent subsurface germination weeds from becoming established before Garden Weed Preventer is activated.

The following weeds can be controlled with proper use of this product: Barnyardgrass, browntop panicum, carpetweed, chickweed, crabgrass (smooth and large), fleabane, Florida pusley, foxtail (green and yellow), goosegrass, ground cherry, johnsongrass seedlings, lambsquarter, lovegrass, nodding spurge, pigweed (red root), purslane and witchgrass.

BEST AVAILABLE COPY

For best results: Sprinkler irrigation should be used when there is no rain within three days after application.

REPLANTING: Replanting crops other than those included on this label in Dacthal treated soil within eight months of application may result in crop injury. If replanting is required because of early crop failure, the planting of onions, seeded melons, potatoes, tomatoes, eggplants, or peppers at this time may result in crop injury. However, all crops on this label may be planted following harvest of a Dacthal treated crop.

Under unfavorable growth conditions, slight crop injury and reduction in yield may result.

<u>CROPS</u>	<u>WHEN TO APPLY</u>	<u>REMARKS</u>
Broccoli, brussel sprouts, cauliflower, cabbage, garlic, collards, kale, mustard greens, and onions	At seeding or transplanting	Apply uniformly to the soil at the rate of 1 lb. per 250 sq. ft. If weeds have emerged, soil should be clean cultivated.
Snap beans, southern peas, (blackeyed)	At seeding	Apply uniformly to the soil at the rate of 1 lb. per 250 sq. ft. at time of planting. Soil should be clean cultivated or weeded prior to application.
Potatoes (whole or cut pieces)	At planting or drag off	Apply uniformly to the soil at the rate of 1 lb. per 250 sq. ft. at planting or following planting. If the top of the beds are to be dragged off, Garden Weed Preventer should be applied after drag off. Soil should be weeded prior to application. Dacthal is not recommended for use on potatoes in California.
Sweet potatoes, yams	At transplanting and anytime up to 6 weeks after transplanting	Apply uniformly to soil at the rate of 1 lb. per 250 sq. ft. at transplanting time. Soil should be weeded prior to application.

BEST AVAILABLE COPY

Tomatoes, eggplant,
and peppers

4 to 6 weeks
after trans-
planting

Uniform application to the
soil should be confined to
a period of 4 to 6 weeks
after transplanting.
Plants should be well
established and growing
conditions favorable for
good plant growth. Soil
should be weeded prior to
application.

Strawberries - new
plantings

At transplanting

Apply uniformly to the soil
at the rate of 1 lb. per
250 sq. ft. at
transplanting time. Soil
should be weeded prior to
application.

Strawberries -
established plantings

Fall and
early Spring

Apply uniformly at the rate
of 1 lb. per 250 sq. ft.
DO NOT apply from first
bloom through harvest.
Application may be made
directly over the plants
without injury.

Nursery stock,
ornamentals and flower
gardens

Early Spring

Apply uniformly to soil at
the rate of 1 pound per 250
sq. ft. Soil should be
weeded prior to application
Delayed or secondary
applications can be made at
any time provided beds have
been cultivated and are
free of weeds.

BEST AVAILABLE COPY

GARDEN WEED PREVENTER CAN BE USED WITH THE FOLLOWING:

Abelia	Ageratum	Alyssum	Andrandea, Jap
Arborvitae	Ash	Aster	Azalea
Babysbreath	Barberry	Bellflower	Birch
Bleedingheart	Bloodleaf	Boxwood	Bugloss
Camellia	Candytuft	Chestnut	Chrysanthemum
Cinquefoil	Coleus	Columbine	Coneflower, Purple
Coreopsis	Coralbells	Cosmos	Cotaneaster
Cottonwood	Crabapple	Cuphea	Cypress
Dahlia	Delphinium	Deutzia	Dogwood
Elaeagnus	Elm	Euonymus	Evening Primrose
Feverfew	Fir	Forget-me-not	Forsythia
Four-O'Clock	Foxglove	Gaillardia	Geranium
Gladiolus	Golddust	Goldentuft	Gum
Heath Holly	Hydrangea	Honeysuckle	Iris
Ivy, Baltic	Ivy, Boston	Ivy, English	Ivy, Wilson
Juniper, Lantana	Larkspur, candle	Lavendar cotton	Lilac
Lily	Locust	Lupine	Magnolia
Maple	Marquerite, golden	Marigold	Mockorange
Morningglory	Moss rose	Mother-of-Thyme	Mountain Laurel
Mourning Bride	Nasturtium	Oak	Orpine
Pachistima	Pachysandra	Peony	Petunia
Pittosporum	Podocarpus	Pocker plant	Poplar
Privet	Privet, variegated	Redbud	Rhododendron
Rose	Sage, scarlet	Snapdragon	Spiderwort, Va
Spirea	Spruce	Stonecrop	Strawflower
Sun drops	Sunflower	Sweet pea	Sycamore
Tree peony	Tulip tree	Viburnum	Violet
Walnut	Weigela	Willow	Warmwood
Yarrow, fernleaf	Yew	Zinnia	

DO NOT USE ON THE FOLLOWING PLANTS:

Buglewood	Button pink	Carnation	Germander
Geum	Mesembryanthemum	Pansy	Phlox
Telanthera	Sweet William		

STORAGE AND DISPOSAL ~ 12 pt.

Do not contaminate water, food or feed by storage and disposal. Keep container closed and away from food, feedstuffs and domestic water supplies. **PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash. **CONTAINER DISPOSAL:** Do not reuse bag. Discard bag in trash.

BEST AVAILABLE COPY

WARRANTY

MANUFACTURER warrants that this product (1) conforms to the ingredient statement on the label and (2) is reasonably fit for the purposes set forth in the Directions For Use. EXCEPT AS SO WARRANTED, THE PRODUCT IS SOLD AS IS. MANUFACTURER MAKES NO OTHER WARRANTY EXPRESS OR IMPLIED.

The manufacturer's directions regarding use of this product are based upon tests believed to be reliable. All statements made concerning this product apply only when used as directed, under normal use conditions. FOLLOW DIRECTIONS CAREFULLY. Timing and method of application, weather and turf conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the manufacturer.

Manufactured by:

WEGRO
P.O. BOX 119
MAUMEE, OH 43537

:5815-35

~~10/28/88~~

5/20/90

BEST AVAILABLE COPY