


#### DIRECTIONS FOR USE

Stir or shake thoroughly to mix contents. Wood should be dry and free from bark to allow best penetration. Soak wood for a minimum of three minutes or apply, liberally, two or more brush coats. All possible surfaces should be covered, especially end grains, joints, and those to be in contact with the soil. Fence posts should be soaked in preservative for 12 to 48 hours.

Net Contents: One Gallon

#### CONDITION OF SALE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



A ready-to-use, black, wood preservative. Use to control destruction of wood caused by rot, termites, or other boring insects. Recommended for exterior wood such as fence posts, timbers, posts and beams which are not to be painted. Multiplies the normal life of wood with an easy, quick, and low cost treatment.

**WARNING:** POISON, FLAMMABLE. Harmful if swallowed, inhaled, or absorbed through skin. May cause skin irritation. Do not breathe vapor or spray mist. Do not get in eyes, on skin, or on clothing. Wash thoroughly after handling. Do not reuse container. Destroy when empty. Keep away from heat and open flame. Use in well ventilated areas. Contains flammable solvents.

3-24-75

2705-330

**CAUTION**

Keep Out of Reach of Children

**MAY CAUSE IRRITATION OF NOSE, THROAT, AND SKIN.**

**AVOID SKIN CONTACT.**

**DO NOT BREATHE DUST.**

**Observe the Following Use Precautions**

Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated. Sulphur may burn foliage when temperatures are high. Do not apply sulphur when shade temperatures exceed or are likely to exceed 90° Fahrenheit.

**DIRECTIONS FOR USE**

This dust should be applied when the air is calm. To obtain the most effective results a good hand or power dusting machine should be used to provide a complete coverage of plants with a uniform coating of dust.

Completely empty contents and bury the unused chemicals at least 18 inches deep in an isolated location away from water supplies. Burn empty container immediately. Exercise extreme caution and stay well away from the smoke. Bury ashes and residues.



**3.4% COPPER—86%  
SULPHUR DUST**

**AN AGRICULTURAL FUNGICIDE  
USE ONLY AS RECOMMENDED!**

**CAUTION KEEP OUT OF REACH OF CHILDREN.  
SEE OTHER CAUTION STATEMENTS ON SIDE PANEL.**

**ACTIVE INGREDIENTS: BY WEIGHT**

Copper expressed as metallic (from Copper Oxide)	3.4%
Sulphur as elemental	86.0%
<b>INERT INGREDIENTS</b>	<b>10.6%</b>
<b>TOTAL</b>	<b>100.0%</b>

**SULPHUR MESH TEST: 93% Passing 325 U.S.  
Sieve.**

**RECOMMENDATIONS**

**PEANUTS:** Peanut Leaf Spot—Apply dust about 60 days after planting, or as soon as spots appear on the leaf.

First Dust—Apply 15 to 20 pounds per acre.

Later Dust—Apply 20 to 30 pounds per acre.

A minimum of three to four applications should be made at two-week intervals. Consult State Agricultural Experiment Stations or State Agricultural Service for complete details on the peanut dusting program best suited to your local conditions.

**NOTICE OF WARRANTY**

Helena Chemical Company warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions for Use and Conditions of Sale, subject to the inherent risks referred to therein.

Helena Chemical Company makes no other expressed or implied warranty, including any other expressed or implied warranty of **FITNESS** or of **MERCHANTABILITY**, and no agent of Helena Chemical Company is authorized to do so except in writing, with a specific reference to this warranty. Any damages arising from a breach of this warranty shall be limited to direct damages, and not include consequential commercial damages such as loss of profit or value etc.