NOV 21 1980

656 185,791

Mr. Robert E. Windler Windler Pest Control, Inc. 116 East 5th Street Fowler, IN 47944

Dear Mr. Windler:

Subject: Windler Rat and House Killer
EPA Registration No. 5836-3
Your Application Dated October 20, 1986

The amendment referred to above, submitted in connection with registration under section 3(c)(7)(A) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable provided that you:

- Submit/cite all data required for registration/reregistration
 of your product under FIFRA section 3(c)(5) when the Agency
 requires all registrants of similar products to submit such data.
- 2. Hake the labeling change listed below before you release the product for shipment bearing the amended labeling:
 - Relocate the Child Hazard warning (Keep Out of Reach of Childrer) to above the Signal Word (Caution).
- 3. Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

85295:Palmateer:CBI-J(b):KENCO:11/12/86:11/19/86:de:vo

		* **		•			
				CONCURRENCES			
SYMBOL >					<u> </u>		
SURNAME	,						
DATE		. ,					
EPA form 1	320-1 (4-81)	1.	<u> </u>		<u> </u>	 OFFI	CIAL FILE COPY

In the future please indicate your Registration Number on your letter when making a submission.

Sincerely yours

William H. Miller Product Hanager (16)

Insetticide-Rodenticide Branch'
Registration Division (TS-767C)

LAKUE ELLIYUM I BILE

WINDLER

RAT AND MOUSE KILLER

READY TO USE - NO MIXING REQUIRED

PELLETED

NET WT. 5 LBS.



WINDLER PEST CONTROL, INC.

Fowler, Indiana CAUTION

Keep Out of Reach of Children See antidate and other Warnings on back panel

ACTIVE INGREDIENT:

(Warfarin) 3-(alpha-acetonylbenzyl) 4-hydroxycoumarin* 0.025% INERT INGREDIENTS 99.975%

TOTAL 100.00%

ACCEPTED with COMMENTS in EPA Letter Dated:

NOV 2 1 1956

Under the "	∴¦e,
Funcicide	7,43
as amenara,	:34
registered une distriction	N. 19
3B36-3	

SPECIAL NOTICE - READ BEFORE USING

Use of this product for selective killing of rats requires controlled isolation of its effects at user's risk. Carefully avoid all described and related dangers and any practice not specifically recommended. Recovery on manufacturer's or seller's civil liability for any cause is hereby expressly limited to refund of purchase price.

Do Not Reuse Container, Destroy When Empty.

FRA. Reg. No. 5836-13

PRECAUTIONARY STATEMENTS: HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN - If ingested, administer Vitamin K1 intramuscrularly or orally, as indicated in bishydroxycoumarin overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD:

Keep Out of Lakes, Streams or Ponds

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS - For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contains that food.

SELECTION OF TREATMENT AREAS - Determine areas where rate aedior mice will most likely find and consume the bait. Generally, these areas are along walls, by graved openings, in or beside burrows, in corners and concealed places between floors and walls, or in locations where rouents or their signs have been observed. Protect bait firth rain or snow. Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain arruminterrupted subbly of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 Jays.

RATS & MICE: Replace contaminated or applied bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container only in a dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse bag. Discard in trash.

· Desta Durcha- ause ie Lunu-



Use of this product for selective killing of rats requires controlled isolation of its effects at user's risk. Carefully avoid all described and related dangers and any practice not specifically recommended. Recovery on manufacturer's or seller's civil liability for any cause is hereby expressly limited to refund of purchase price.

Do Not Reuse Container. Destroy When Empty.

PRECAUTIONARY STATEMENTS: HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN - If ingested, administer Vitamin K1 intramuscrularly or orally, as indicated in bishydroxycoumaria overdoses. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD:

Keep Out of Lakes, Streams or Ponds

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS - For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamper-proof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS - Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walfs, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walfs, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger-placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain as unintegrupted supply of fresh bait for at least 15 days.

RATS.& MICE: Replace contaminated or spoiled hait immediately. Collect and dispose of all dead crimals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and restensh bait as needed.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

STORAGE: Store in original container only in a dry place inaccessible to children and pets.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Do not reuse bag. Discard in trash.