

PR 16

REG 5857-4

RESTRICTED USE PESTICIDE

For retail sale to and use only by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's certification.



FOR USE AGAINST LISTED INSECTS WHICH INFEST STORED COMMODITIES, SPECIFIED PROCESSED FOODS, AND ANIMAL FEEDS

* Active Ingredient, Aluminum phosphide 55% Inert Ingredient: 45%
Pat No 3132067 and 3866347 E.P.A. Reg. No. 5857-4-AA



POISON



DANGER Keep out of reach of children

See antidote and other precautions on back panel

Contents: 4 packages of 165 pellets each. Net wt. 386 gms. (.87 lbs.)

Manufactured by **DEGESCH GMBH · FRANKFURT AM MAIN**

Federal Republic of Germany
EPA Est. 34889-WQ-1

Distributed by **PHOSTOXIN SALES, INC.**

Box 468, Alhambra, Cal. 91802 · Telephone (213) 283-2781

DISPOSAL OF CONTAINERS:

Decontaminate first by filling them with water after the PREPACS have been dispensed



POISON



ANTIDOTE:

Remove victim to open air and remove any contaminated clothing. If ingested, give 2 tablespoons of salt in a glass of warm water to induce vomiting. Repeat until vomit fluid is clear. Call a physician immediately.

DANGER:

PHOSTOXIN releases poisonous gas when moist. Do not breathe dust or fumes. Poisonous if swallowed. Do not get in eyes or on skin. KEEP AWAY FROM LIQUID OR WATER. Wear gloves when handling pellets. PHOSTOXIN to be used only by experienced adult personnel. Do not open containers except for immediate usage. Always open to open air. Store only in cool, dry, locked, ventilated room. Protect from moisture. open flames or heat. Keep available a gas-mask and respirator approved by the U. S. Department of Interior, Bureau of Mines, for atmospheric protection.

SYMPTOMS:

Pressing sensation in the chest, nausea, and dizziness

FOR FUMIGATION OF LISTED RAW AGRICULTURAL COMMODITIES, SPECIFIED PROCESSED FOODS, AND ANIMAL FEEDS:

For space treatment of stored products insects in stored barley, corn, millet, oats, peanuts, rice, rye, softener seeds, seed and grain vegetables (except soybeans), sorghum, sunflower seeds, wheat, specified processed foods, and animal feeds. These commodities may be fumigated under gas-proof tarpaulins in sealed box or hopper cars (plastic or rubber) or other suitable enclosures. The package shall be fastened to a substantial support and it shall not come in contact with the raw commodity, specified processed food, or animal feed. After fumigation, the pellets dust remaining shall be dropped off by burying entire package or by immersing it in water to which a small amount of washing agent has been added. Do not dispose of package in toilet.

BARGE FUMIGATION IS REGULATED BY U.S. COAST GUARD REGULATION 48CFR 147A

Unmanned barges are fumigated as follows:

For space treatment of listed raw commodities, specified processed foods, and animal feeds, the package of pellets shall be fastened to a substantial support and it shall not come in contact with the raw commodity, specified processed food, or animal feed.

Following application of the fumigant, the barges must be sealed and warning placards attached.

Persons unloading of unmanned barges make appropriate tests to ascertain safety of cargo area.

Dosage for all listed raw commodities, specified processed foods and animal feeds is one package per 1000-1800 cubic feet of storage space. DO NOT SUBDIVIDE PACKAGE.

SEE ENCLOSED SHEET FOR THOSE LISTED PROCESSED FOODS WHICH MAY BE FUMIGATED.

The times of exposure required are approximated in the following guide:

at 54-59°F (12-15°C) - 4 days are required
at 60-68°F (16-20°C) - 3 days are required
at more than 69°F (20°C) - 2 1/2 days, not less

Do not fumigate when product is below 40°F (5°C)

Average finished food 48 hours before it will be offered to the consumer.

PHOSTOXIN has been found effective against listed grain insects and their pre-adult stages - that is, eggs, larvae, and pupae. Grain and rice weevils, saw-toothed grain beetle, flat or rusty grain beetle, lesser grain borer, nut-weevil, flour beetle, Cadelle dried fruit moth, Mustier green flour moth, Indian meal moth, Angoumois grain moth.

Tobacco may be fumigated with PHOSTOXIN at a dosage rate of one package per 1000 cubic feet. Aspartin lime on hogheads will be not less than 3 days, all others 2 days.

Under no condition shall any processed food or animal feed come in contact with PHOSTOXIN or with PHOSTOXIN residue. PHOSTOXIN shall not be placed in or attached to packages intended for consumers or articles delivered to retailers.

Warranty: Seller warrants that the product conforms to its commercial description and when used according to label directions under normal conditions of use, it is reasonably fit for the purposes stated on the label. Seller makes no other warranty, either express or implied, and Buyer assumes a risk should the product be used contrary to label instructions.

4/74

ACCEPTED

OCT 1 1979

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 5857-4

BEST AVAILABLE COPY

Handwritten mark resembling '11' or '1/1'.