

TEXO



orporation
CINCINNATI, OHIO 45212

ACTIVE INGREDIENTS

Alkyd 50% C14 40% C12 10% C10
100% active ingredient

INERT INGREDIENTS

Water

Surfactant
Stabilizer
Colorant

STORAGE AND DISPOSAL

Store in original container
Do not use if container is damaged
Do not use if contents are empty

Keep out of reach of children
Do not use if contents are empty
Do not use if container is damaged

Do not use if contents are empty
Do not use if container is damaged
Do not use if contents are empty

SEPTODOR

Disinfectant-Sanitizer-Deodorizer-Algaecide-Slime

Do not use in a manner inconsistent with the label. Do not use in food processing plants and in manufacturing plants where a disinfectant is required.

If surfaces have been soiled, they can be treated with Septodor.

For disinfecting walls, floors, and similar hard surfaces apply a solution of 1 part Septodor to 100 parts of water.

For disinfecting floors, apply a solution of 1 part Septodor to 100 parts of water.


For disinfecting walls, floors, and similar hard surfaces apply a solution of 1 part Septodor to 100 parts of water.

For disinfecting walls, floors, and similar hard surfaces apply a solution of 1 part Septodor to 100 parts of water.

For disinfecting walls, floors, and similar hard surfaces apply a solution of 1 part Septodor to 100 parts of water.

For disinfecting walls, floors, and similar hard surfaces apply a solution of 1 part Septodor to 100 parts of water.

FOR INDUSTRIAL AND COMMERCIAL USE



Hazards To Humans and Domestic Animals
DANGER: Keep Out of Reach of Children.

Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.

FIRST AID: In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated

clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Environmental Hazards: This pesticide is toxic to fish. Do not discharge into lakes, streams, ponds or public waters unless in accordance with a NPDES permit. For guidance, contact your Regional Office of the Environmental Protection Agency.

SEPTODOR

Disinfectant-Sanitizer-Deodorizer-Algaecide-Slimicide

This is a violation of federal law to use this product in a manner not intended with the label. Do not use in Hotels, Motels, Food Service, or in Medical Facilities. Do not use in swimming pools where a disinfectant, sanitizer, deodorizer, or algaecide is required.

After surfaces have been cleaned, the following instructions apply to the use of SEPTODOR as follows:

- For disinfecting walls, floors, and similar surfaces, use a solution of 100 ppm (one ounce per gallon) of water.
- For sanitizing walls, floors, and similar surfaces, use a solution of 50 ppm (one-half ounce per gallon) of water.
- For deodorizing walls, floors, and similar surfaces, use a solution of 25 ppm (one-quarter ounce per gallon) of water.

For more information, contact your local distributor or write to:

SEPTODOR

| Phenol Coefficients - AOAC Method | |
|-----------------------------------|----|
| Salmonella Typhosa | 50 |
| Staphylococcus Aureus | 75 |

FOR USE AS COOLING TOWER ALGAEICIDE SLIMICIDE

PRE-TREATMENT: Add this product to the surface of the cooling tower.

INITIAL DOSE: In a clean system with no signs of microbial slime growth visible, add six gallons of SEPTODOR per 10,000 gallons of cooling water. If algae and microbial growth are visible, add six gallons of SEPTODOR per 10,000 gallons of cooling water.

SEQUENT TREATMENT: To work with the above Field Equipment, After the initial dose, maintain a concentration of 10 ppm of the active ingredient in the water by feeding SEPTODOR at a rate of one gallon per 10,000 gallons of cooling water. If microbial growth with persistent slime is observed, add three gallons of SEPTODOR per 10,000 gallons of cooling water as a slug dose. This slug dose should be followed with the maintenance concentration of 10 ppm active ingredient.

INDUSTRIAL AND COMMERCIAL USE ONLY