

## FOGICIDE

SPACE SPRAY INSECTICIDE


A high level, synergized pyrethrin type insecticide concentrate for space application by fogging and spraying machines and similar equipment. FOGICIDE can be used around food, food products, utensils and equipment provided that no exposed food is present and provided that food utensils and surfaces that will be in contact with food are covered and provided that fogging and spraying is not performed in food processing plants while plant is in operation.

Properly applied, FOGICIDE kills the accessible exposed stages of flies, ants, mosquitos, boxelder bugs, silverfish, flying moths, gnats, crickets, roaches, spiders, scorpions, wasps, hornets, clover mites, cheese mites, cheese skipper, and most common stored-grain pests, including granary weevil, rice weevil, grain beetle, spider beetle, angoumois grain moth, Mediterranean flour moth and grain mites.

ctive INGREDIENTS	. 100.0	%
*Piperonyl Butoxide, Technical	1.0	%
Pyrethrins	0.10	)%
Petroleum Oil	98.90	)%
*Equivalent to 0.8% of (Butylcarbityl)		
(6-propylpiperonyl) ether and to 0.2%		į

**USDA Registration No. 5870-11** 

related compounds.



FOR INDUSTRIAL AND COMMERCIAL U

#### **DIRECTIONS**

#### Flying Insects

FLIES, MOSQUITOES, GNATS: Direct the spray toward the upper portions of the enclosure and fill the atmosphere with mist, using 2 to 3 ounces of the product to each 1,000 cubic feet of space. If possible, keep the doors and windows closed for 10 minutes after applying the spray.

#### Crawling Insects (Exposed or Accessible Stages)

ROACHES, SILVERFISH, SPIDERS: Spray infected areas thoroughly and direct the liquid behind baseboards, moldings, cabinets, shelves, sinks, door and window frames and pipes, toward drains and dark corners, and into cracks, crevices, recesses and similar hiding places. peat as needed.

ANTS: Treat shelves, cabinets and storage areas. Spray ant runs, trails, baseboards, thresholds and other places of entry, and treat nests liberally.

FLEAS, TICKS: Direct the spray toward lower wall portions, baseboards, rugs, carpets and floors of infested areas. Spray animal quarters, kennels and the bedding of pets and other domestic animals.

BEDBUGS: Apply liberally to top, ends, sides and bottom of mattresses, spraying all tufts, seams and folds. Spray springs and frame, directing the liquid into all cracks, crevices and lock joints. Spray walls, baseboards, molding, floors, closets and bedroom furniture with special attention to protected bedbug hiding and breeding places.

CLOTHES MOTHS AND CARPET BEETLES: Remove infested articles from storage, brush thoroughly and, if possible, air several hours in direct sunlight. Apply spray liberally to all surfaces of empty chests, closets, bureaus and storages, directing the liquid into cracks, joints and crevices.

### PYROKILL INSECT SPRA

37-73

#### CAUTION

KEEP OUT OF REACH OF CHILDREN

See Side Panel for Required Causion

ACTIVE INGREDIENTS	·
Petroleum Distillate	<b>98</b>
Piperonyl Butoxide*	1
Pyrethrins	0

\*Equivalent to 0.8% of (butylcarbitol) (6-propylpiperonyl) ether and of related compounds.

EPA No. 3837-32

Manufactured by:

#### LUBAR COMPANY

1708 Campbell Street

Kansas City, Miss



# FOGICIDE SPACE SPRAY INSECTICIDE

Property at all FOGICIDE kills the accessible exposed stances of ties, and amosquitos bouldider bugs, silverfish, flying meths ghats, crickets roaches, spiders, scorpions, wasps, homets diever mites, cheese mites, cheese skipper, and most common stored-grain pests including granary weevil, rice weevil, grain beetle, spider beetle, angoumois grain moth. Mediterrate in four moth and grain mites.

Active INGREDIENTS

- \*Piperonyl Butoxide Technical
- Pyrethmas
- Petroleum Oil
- \*Equivalent to 0.8% of (Butylcarbity) (6-propylpiperonyl) ether and to 0.2%

related compounds.

**USDA Registration No. 5870-11** 



#### CAUTION: READ BEFORE USING

#### HARMFUL IF SWALLOWED KEEP OUT OF REACH OF CHILDREN

Do not remain in treated areas and ventilate the area after treatment is completed. FOGICIDE is an oil base material and should not be sprayed, stored or otherwise exposed to an open flame.

In order to avoid any possible explosive hazard, not more than 1 gallon should be sprayed per 40,000 cu. ft. of enclosed space.

Remove or cover all exposed foods or foodstuffs. Cover all utensils and surfaces that will be in contact with food. Do not spray or fog in food processing plant while plant is in operation.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in safe place.

The contract was translated as important that the traction of the

0.3 of the distributions of 0.00 c. The 0.3 of 0.3 of 0.3

No DC in the trying insects, if possible even a recommendation to

thristes after application