5 POUNDS NET WT.

# E-Z-KIL

Contains 9.50% Proline Ama

# KILLER BAIT

# Ready to use-No mixing required

31 TIVE INGREDIENTS

Was faring 3 (a-west explorency of Hydroxycountering) (2 quinoxaling, sulfacilité de Sulfaquinoxaline 0.02'). ART INGREDIS NIS 9945 7115 100 (1600)

#### SATISFACTION GUARANTEED

If used according to directions, E-Z Kil is guaranteed to give satisfactory control of rats and mice. Continued use of E-Z Kill will keep these pests under control. Return this package and unused portion to the manufacturer for a refund of purchase price if E-Z Kil does not give you satisfaction.

## Ready to use-No mixing required

TIVE INGREDIENTS.

arka na h ta-acet arthenzy. A hyd. ayeonuna . — 1913 - Pequinoxaliry sulfarilandder Sulfamuinoxaliae (1.025), PERT INGREUMENTS

TO TAL


100,000%

#### SATISFACTION GUARANTEED

If used according to directions, E-Z Kil is guaranteed to give satisfactory control of rats and mice. Continued use of E-Z Kill will keep these pests under control. Return this package and unused portion to the manufacturer for a refund of purchase price if E-Z Kil does not give you satisfaction.

"Prolin is a registered trademark of the Wisconsin Alumni Research Foundation for a rodenticide which is manufactured under license from the Foundation. U.S. Patent Nos. 2,427,578 and 3,113,071."


### **CAUTION**



**Keep Out Of Reach Of Children** 

See antidote and other warning on back panel

U.S.D.A. Reg. No. 5882-1



Contaminated or spoiled baits should be replaced promptly.

Baiting should continue until all signs of feeding have stopped.

"Where a continuous source of infestation is present (nearby dumps) permanent bait stations should be established and replenished as needed."

1-22-71

#### DIRECTIONS FOR RAT CONTROLL

Four ounces of bait per station should be the minimum amount used.

Bait should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes.

An uninterrupted supply of bait should be maintained for at least 10 days.

#### DIRECTIONS FOR MOUSE CONTROL

Tablespoon amounts (1/4 to 1/2 ounce) of bait should be placed at 8-12 foot intervals.

Bait should be placed in locations not accessible to children; pets, and domestic animals, or in tamper-proof bait boxes.

An uninterrupted supply of bait should be maintained for at least 15 days.

#### DIRECTIONS FOR RAT CONTROLL

Four ounces of bait per station should be the minimum amount used.

5882-1

Bait should be placed in locations not accessible to children, pets, and domestic animals, or in tamper-proof bait boxes.

An uninterrupted supply of bait should be maintained for at least 10 days.

#### DIRECTIONS FOR MOUSE CONTROL

Tablespoon amounts (1/4 to 1/2 ounce) of bait should be placed at 8-12 foot intervals.

Bait should be placed in locations not accessible to children; pets, and domestic animals, or in tamper-proof bait boxes.

An uninterrupted supply of bait should be maintained for at least 15 days.

#### CAUTION

This product contains as its active ingredient an anti-coagulant chemical (Warfarin) which if taken by humans, domestic animals, or pets, may reduce the clotting ability of the blood and internal hemorrhage may result. PROTECT FEEDING PANS FROM CHILDREN, DOGS, CATS, AND LIVESTOCK. In case baits are eaten, give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Call a physician immediately.

NOTE FOR PHYSICIANS: When a human has been known to have accidentally ingested this product, blood transfusions combined with intravenous injections and oral doses of Vitamin K are indicated as in the case of hemorrhage caused by overdoses of Dicumarol. (Reg.)

#### **NOTICE**

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.