


Black Leaf
Chlordane Dust



ACTIVE INGREDIENT:
 Technical Chlordane 10%
INERT INGREDIENTS 90%
 Total 100%
 Equivalent to 6% Octachloro-4,
 7-Methanotetrahydroindane and
 4% related compounds.

**NET WEIGHT
 ONE POUND**

**CAUTION
 KEEP OUT OF THE
 REACH OF CHILDREN**
 See side panel for other cautions

**for ants and
 lawn insects**

KILLS *ants,
 cutworms, wireworms
 japanese beetle larvae
 lawn moths*

CAUTION: Avoid inhalation. Do not contaminate food, feed and drinking water.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry. Toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be harmed. Do not contaminate water by cleaning equipment, or disposal of wastes.

ANTIDOTE

External—in case of spillage on skin, wash promptly with soap and water. Internal—Emetic of 1 tablespoonful mustard in tumbler of water. Call a physician.

SELF-APPLICATOR PACK: Perforate or punch holes in top of package making sure that plug in top of can is firmly in place. Shake out dust in package through scientifically measured holes for even application.

HAND DUSTER APPLICATION: It is not necessary to perforate holes in top of package. Simply remove plug in top of can and empty desired amount into your duster. Replace plug for continued storage.

CONTAINER DISPOSAL: Do not reuse empty container. Rinse equipment and containers and dispose of wastes by burying in non-croplands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning. Stay away from smoke or fumes.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**Chloro
 Dus**

GENERAL DIRECTIONS

DIRECTIONS FOR USE OF BLACK LEAF
 Please read directions carefully. of this product. Keep children and pets off treated areas until washed into the soil and grass is dry.

USE	INSECTS CONTROLLED	
Lawn and Turf	Ants Chiggers Fire Ants Japanese Beetle Larvae	Lawn Sod Wireworms
Vegetables	Armyworms Cabbage Maggot Cabbage Worms Cucumbers Lettuce Tomato	Greenhouse Bugs Harlequin Serpentine Leafhoppers Squash Stinkbugs Wireworms Flea Beetles
Outdoors	Ants, Boxelder Beetles	

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

BLACK LEAF
 DISTRICT
 M ELGIN, ILL.

CAUTION: Avoid inhalation. Do not contaminate food, feed and drinking water.

Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

Toxic to fish and wildlife. Keep out of any body of water. Birds feeding on treated areas may be harmed. Do not contaminate water by cleaning equipment, or disposal of wastes.

ANTIDOTE

External—in case of spillage on skin, wash promptly with soap and water. **Internal**—Emetic of 1 tablespoonful mustard in tumbler of water. Call a physician.

SELF-APPLICATOR PACK: Perforate or punch holes in top of package making sure that plug in top of can is firmly in place. Shake out dust in package through scientifically measured holes for even application.

HAND DUSTER APPLICATION: It is not necessary to perforate holes in top of package. Simply remove plug in top of can and empty desired amount into your duster. Replace plug for continued storage.

CONTAINER DISPOSAL: Do not re-use empty container. Rinse equipment and containers and dispose of wastes by burying in non-croplands away from water supplies. Containers should be disposed of by punching holes in them and burying with wastes or by burning. Stay away from smoke or fumes.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Chlordane Dust

GENERAL DIRECTIONS FOR USE

DIRECTIONS FOR USE OF BLACK LEAF 10% CHLORDANE DUST
Please read directions carefully. Consult chart for the many uses of this product.
Keep children and pets off treated areas until this material is washed into the soil and grass to dry.

USE	INSECTS CONTROLLED	DOSAGE AND APPLICATION
Lawn and Turf	Ants Chiggers Fire Ants Japanese Beetle Larvae Lawn Moths Sod Webworms Wireworms	Apply at the rate of 1-lb. per 700 sq. feet of area. Wash into soil by sprinkling with water.
Vegetables Cabbage Broccoli Cauliflower Celery Cucumbers Lettuce Tomato	Armyworms Cabbage Maggot Cabbage Worms Cucumber Beetles Cutworms Flea Beetles Green Stink Bugs Harlequin Bug Serpentine-Leafminer Squash Bugs Stink Bugs Wireworms	Apply at the rate of ¼-lb per 700 sq. feet. Dust around base of plants preferably in early morning when air is still. Repeat applications at 10 day intervals. Do not apply after edible parts begin to form.
Outdoors	Ants, Boxelder Bugs, Wasps	Dust nests, runways and around tree trunks, and lower limbs of Boxelder Trees. Dust entrances.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

DISTRIBUTED BY
BLACK LEAF PRODUCTS CO.
M ELGIN, ILLINOIS 60120
00-903

