



# Fruit Tree Spray

MAR. 10, 1982

5887-16

## Protect your fruit trees

**Contains: methoxychlor, malathion, captan. A complete insecticide, fungicide and miticide for insect and disease control on fruit trees.**


ACTIVE INGREDIENTS	BY WT.
Technical Methoxychlor*	10.0%
Captan — N-(trichloromethyl)- iso-4-cyclohexene-1,2-dicarboximide	7.0%
Malathion—O,O-dimethyl Dithiophosphate of diethyl mercaptoacetate	5.0%
TOTAL	72.0%
INERT INGREDIENTS	100.0%

\*Equivalent to 8.1% 2,2-bis(p-methoxyphenyl) 1,1-trichloroethane and 1.2% chloro isomer and related compounds.

**KEEP OUT OF  
REACH OF CHILDREN**

### CAUTION

See rear panel for statement of practical treatment and other precautionary statements



EPA EST. NO. 5887-IL-1  
EPA REG. NO. 5887-16  
Net Weight 16 oz. (1 pound)

## Fruit Tree Spray

### STATEMENT OF PRACTICAL TREATMENT

If In Eyes: Flush with plenty of water. Get medical attention if irritation persists.  
If Swallowed: Drink a glass of water and induce vomiting by touching finger to back of throat. Get medical attention.  
If On Skin: Wash with plenty of soap and water. Remove and wash contaminated clothing.

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

Harmful If swallowed. Avoid inhalation of dust or spray mist. Avoid contact with skin. Do not get in eyes, on skin, or on clothing. Use with adequate ventilation. Avoid contamination of food and feedstuffs.

#### ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. This product is highly toxic to bees exposed to direct treatment or residues in crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**SPRAY APPLICATION:** One pound of this material will make 10 gallons of fruit tree spray. For smaller quantities, use 5 level tablespoons to each gallon of water. Mix well. Agitate solution while spraying. (BLACK LEAF FRUIT TREE SPRAY may be applied as a dust by use of an efficient blower and directing the dust to all portions of the tree being treated.)

#### AMOUNT OF SPRAY NEEDED PER TREE

SIZE	DILUTED SPRAY
Young or Dwarf	Approx. 1 to 2 pints
Medium	Approx. 1 to 2 quarts

**APPLES, PEARS:** To control scale, brooks fruit spot, woolly and green apple aphids, pacific clover and red spider mites, codling moth, curculio, apple maggot, Japanese beetle, and bitter rot, apply as follows:

1. When blossom stems separate and before blossoms open. 2. When 75% of the petals have fallen. 3. 10 days later. 4. 10 days to 2 weeks later. 5. Continue at 2-week intervals to 21 days before harvest.

**PEACHES:** To control brown rot, peach scale, European red mite, red spider, plum curculio, Japanese beetle, European fruit scale, pacific and clover mites, apply as follows:

1. Full bloom. 2. When shucks begin to split. 3. 10 days later. 4. 10 days later. 5. 2 weeks later. 6. Do not apply within 21 days of harvest.

**PLUMS (PRUNES):** To control pacific, European and red spider mites, plum curculio, mealy plum and green peach aphids and Japanese beetle, apply as follows:

1. When trees are in full bloom. 2. When shucks are falling. 3. 10 days later. 4. 2 weeks later. 5. Continue at 2-week intervals if necessary. Do not apply later than 21 days before harvest.

If control of disease on above fruit crops is necessary within 21 days of harvest, apply a wettable sulfur spray. Remove residues by washing, wiping, brushing or other effective means.

#### DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection. This container sold by weight and not by volume. If it does not appear full when opened, this is because contents have settled during shipping and handling.

**NOTICE:** Buyer assumes all risks of use, storage, or handling of this product not in strict accordance with directions given herewith.

**DISTRIBUTED BY**  
**BLACK LEAF PRODUCTS COMPANY**  
ELGIN, ILLINOIS 60120  
60-302