

CAUTION

HUMANS - Wash hands after use. Harmful if swallowed. Keep out of reach of children. Do not use or store hear heat or open flame.
PLANTS — Avoid spraying BLACK LEAF
DORMANT SPRAY on foliage during
high temperatures or when there has
been a lack of rain. Do not apply to broad-leaved evergreens during freezing or near freezing temperatures. Not to be used on Maidenhair Ferns, Cocas

Palms, or Maples.

रिविधाना आसार स्थार lo evole त्र्मार्ग रेमार्ग

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

ACTIVE INGREDIENTS: By Wt. Petroleum Oils 97% (Pet. Oils-Min. U.R. 94% Class-Light-Med.) *Mixture of Polyglycol Esters and fatty amides 100%

DISTRIBUTED BY BLACK LEAF PRODUCTS CO.

> CHICAGO, ILLINOIS U.S.A. USDA REG. NO. 5887-37

sticker-spreader during growing season

AIDS IN CONTROL OF:

Garden Pests

Scale Insects **Aphids Red Spiders**

White Flies Mealybugs

CAUTION KEEP OUT OF THE REACH OF CHILDREN

See side panel for other cautions

GENERAL DIRECTIONS

PLEASE READ OUR LABEL CAREFULLY BEING SURE TO NOTE ALL INSTRUC-TIONS, WARNINGS, AND CAUTIONS.

Shake well before using, Thoroughly wet all plant pests, as well as all portions of the plants being sprayed.

DORMANT SPRAY: BLACK LEAF DORMANT SPRAY will control Red Spiders. Mites, Scale Insects, Mealybugs on Dormant Shrubs, Evergreens, Fruit Trees, and Shade Trees, as well as other woody plants, especially during the dormant season. Mix 5 ozs. of BLACK LEAF DORMANT SPRAY in one gallon of water when preparing for spraying.

SPREADER OR STICKER: BLACK LEAF DORMANT SPRAY can be combined with other Black Leaf Products Sprays, such as BLACK LEAF 40, BLACK LEAF 50% MALATHION SPRAY, BLACK LEAF SHEEN (a fungicide) and will increase the efficiency of the spraying since it will cause the insecticides or fungicides to adhere to the insects or plant surfaces which are being treated. When using as a Spreader-Sticker, the insecticides or fungicides may be used at the regular strength. Add four teaspoonsful of BLACK LEAF DORMANT SPRAY to each gailon of spray being mixed.

SPRAY FOR GROWING SEASON: Scale Insects, Red Spiders, Mites, Mealybugs, White Fly Larvae on Citrus, Fruit, Shade Trees, Evergreens, Camellias, Azaleas, Hardy Shrubs, can be controlled by using five tablespoonsful BLACK LEAF DORMANT SPRAY to each gallon of water. This application may be used with safety during the growing season.

WINTER SEASON: Mix five ounces of BLACK LEAF DORMANT SPRAY with one gallon of water for control of Scale insects and insect eggs that were left on the plants. This will help to control many scale insects. BLACK LEAF DORMANT SPRAY will be effective in destroying larvae of Red Spider

HOUSE PLANT LEAF SHINE: BLACK LEAF DORMANT SPRAY can be used to brighten the leaves of hardy house plants, by using with a mixture of three tablespoonsful of Black Leaf Dormant Spray per gallon of water. (Do not use on African Violets.)

IN THE GREEN HOUSE: Use two tablespoonsful of BLACK LEAF DORMANT SPRAY per gallon of water to control Scale Insects, Red Spider, Mealybugs on Hardy Shrubs, and White Flies, Allow 60 days between sprays.

