

5887-37 101 pm/2

EPA-IL

10-20-80

REC. AVAILABLE COPY

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS

CAUTION

Harmful if swallowed. If swallowed, do not induce vomiting. Avoid breathing of vapors and spray mists. Wash after handling.

ENVIRONMENTAL HAZARDS

Kill fish, lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Please read entire label carefully before using, being sure to follow all instructions and cautions. Shake well before using. Thoroughly wet all plant pests, as well as all portions of plants being sprayed. Shake diluted spray frequently during application.

DORMANT SPRAY BLACK LEAF DORMANT SPRAY will control red spiders, mites, scale insects, mealybugs on dormant shrubs, evergreens, fruit trees and shade trees, as well as other woody plants, especially during the dormant season. Mix 3 ozs. of BLACK LEAF DORMANT SPRAY in one gallon of water when preparing for spraying.

SPREADER-STICKER: BLACK LEAF DORMANT SPRAY can be combined with other BLACK LEAF PRODUCTS SPRAYS such as BLACK LEAF 40, BLACK LEAF 50% MALATHION SPRAY and will increase the efficiency of the spraying since it will cause the insecticides or fungicides to adhere to the insects or plant surfaces which are being treated. When using as a spreader-sticker the insecticides or fungicides may be used at the regular strength. Add four teaspoonsful of BLACK LEAF DORMANT SPRAY to each gallon of spray being mixed. Observe all cautions and limitations on labeling of all products used in mixtures.

SPRAY FOR GROWING SEASON: Scale insects, red spiders, mites, mealybugs, white fly larvae on citrus, fruit, shade trees, evergreens, camellias, azaleas, hardy shrubs, can be controlled by using five tablespoonsful of BLACK LEAF DORMANT SPRAY to each gallon of water. This application may be used with safety during the growing season.

WINTER SEASON: Mix five ounces of BLACK LEAF DORMANT SPRAY with one gallon of water for control of scale insects and insect eggs that were left on the plants. This will help to control many scale insects. BLACK LEAF DORMANT SPRAY will be effective in destroying larvae of red spider and moths.

NOTE ON CITRUS: Do not apply when trees are in or near fruiting. Oil applied during August and September may affect seeds content retard color. Oil applied after October 1 may increase cold damage to trees.

Do not use dormant sprays in combination with or immediately preceding or following applications of Disulfoton compounds, sulfur, Captain, Folpet, Dyrene, Karathane, Morestan, or Sevin. You must determine final to fit local growth and climatic conditions.

HOUSE PLANT LEAF SPRAY: BLACK LEAF DORMANT SPRAY can be used to brighten the leaves of hardy house plants by using a mixture of three tablespoonsful of BLACK LEAF DORMANT SPRAY per gallon of water. (Do not use on African Violets.)

IN THE GREEN HOUSE: Use two tablespoonsful of BLACK LEAF DORMANT SPRAY per gallon of water to control scale insects, red spiders, mealybugs on hardy shrubs and white flies. Allow 60 days between sprays.

STORAGE AND DISPOSAL

Keep oil container tightly closed in storage to prevent entry of water. Do not reuse empty container. Wrap container and put in trash collect in.

NOTICE: Eyer assumes no risks of use, storage or handling of this material not in strict accordance with directions given here with.

DISTRIBUTED BY
BLACK LEAF PRODUCTS CO.
ELGIN, ILLINOIS 60120

66-221




Dormant Spray

Spreader-sticker during growing season

For year round use to control garden pests: aphids, mealybugs, red spiders, scale insects and white flies.

ACTIVE INGREDIENT	BY WEIGHT
Paraffinic Oil	98.8%
INERT INGREDIENTS	1.2%
TOTAL	100.0%
Unsulfonated Residue	92%, min



**KEEP OUT OF
REACH OF CHILDREN**

CAUTION

See side panel for other cautions

EPA EST. NO. 5887-IL-1
EPA REG. NO. 5887-37

Net Contents 32 fl. oz. (1 quart)

