DIRECTIONS FOR LISE:

Mix Black Leaf Maneb rungicide with water, using the level tablespoorful for each gallon of spray desired, if foliage is difficult to wet, add 8 to 10 grops of a spreader-sticker to each gallon. You may also ado most Black Leaf insecticides to this spray mixture, in order to control both fungus as well as other plant pests.

Use a Black Leaf garden sprayer. Opray plants thoroughly applying to bottom and lop surfaces of foliage and to the fruit. Always start these applications before the diseases are apparent and by making applications at 5 to 10 day intervals you will protect your plants through the growing season. Where there is a severe outbreak, shorten the interval between applications. Particularly in the case of black spots on roses, begin applications when the first leaves unford and repeat

TO MAY 1

nine's **, ,** Mer.

MA I SAL to an .

Effective for control of black spot on roses

Effective for control of black spot on roses







at weekly intervals.

Black Leaf Maneb Fungicide can be used to contol fungus on vegetables and grapes as follows:

IOMATOES: Early blight, late blight, Septoria leaf spot, gray leaf spot (Stemphylium) and anthracnose — Do not use on tender young seedlings in greenhouses as injury may occur.

POTATOES: Early blight and late blight

CARROTS: Leaf spot diseases.

CUCUMBERS: Downy mildew, anthracnose.

LIMA BEANS: Downy mildew. (Do not apply within 4 days of harvest

ONIONS: Downy mildew, purple blotch (blight) and blast (Botrytis leaf blight).

SPINACH: Downy mildew (blue mold) and white rust—Do not apply later than 7 days before picking, and remove residues by washing or other effective means. GRAPES: (Great Lakes area) Black rot - Spray just before bloom, just after bloom, and 7 to 10 days later.

LETTUCE AND ENDIVE: Downy mildew - Do not apply within 7 days of picking. Remove residues from head lettuce by stripping and trimming; and from leaf lettuce and endive by washing or other effective means.

CABBAGE: Alternaria leaf spot and downy mildew -- Do not apply within 7 days of picking.

CAUTION

May irritate eyes, nose, throat, and skin. May be harmful if inhaled or swallowed.

Avoid breathing dust or spray mist. Avoid contact with skin, eyes, and clothing. Keep away from fire or sparks. Store away from children, domestic animals, and food stuffs.

In case of contact, flush with plenty of water; for eyes, get medical attention.

NOTICE TO BUYER — SELLER makes no warranty of any kind, express or implied, concerning the use of this product. BUYER assumes all risk of use or handling whether in accordance with directions or not.

!MPORTANT:

KEEP JAR TIGHTLY CLOSED

Effective for control of black spot on roses

Effective for control of black spot on roses

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

See side panel for other cautions USDA REG. NO. 5887-38 Maneb Fungicide controls many plant diseases or fungus on roses--tomatoes --and other garden plants.

ACTIVE INGREDIENT

Manganese ethylenebisdithiocarbamate (Manganese equivalent as metallic 16.5...)
INERT INGREDIENTS

20 O TOTAL 100

80 ୯

This bottle will make 23 gallons of spray 512 oz bottle

BLACK LEAF PRODUCTS COMPANY CHICAGO, ILLINOIS