



Tomato and Vegetable DUST

FUNGICIDE INSECTICIDE

BLACK LEAF TOMATO AND VEGETABLE DUSK

CROPS	DISEASES & PESTS CONTROLLED	AMOUNT TO USE	MADE BY
TOMATOES	Septoria, blight, wilt, mosaic, etc.	1/2 lb. per acre	W. H. H. Co.
POTATOES	Septoria, blight, wilt, mosaic, etc.	1/2 lb. per acre	W. H. H. Co.
PEPPERS	Septoria, blight, wilt, mosaic, etc.	1/2 lb. per acre	W. H. H. Co.
MAIZE BEANS	Septoria, blight, wilt, mosaic, etc.	1/2 lb. per acre	W. H. H. Co.
DRY BEANS	Septoria, blight, wilt, mosaic, etc.	1/2 lb. per acre	W. H. H. Co.
STRING BEANS	Septoria, blight, wilt, mosaic, etc.	1/2 lb. per acre	W. H. H. Co.
SWEET CORN	Septoria, blight, wilt, mosaic, etc.	1/2 lb. per acre	W. H. H. Co.
SQUASH	Septoria, blight, wilt, mosaic, etc.	1/2 lb. per acre	W. H. H. Co.
PUMPKINS	Septoria, blight, wilt, mosaic, etc.	1/2 lb. per acre	W. H. H. Co.

One gallon will cover 50 sq. ft.

Sept 18-1979

5527-70

Please read label carefully and use strictly in accordance with label cautions, warning and directions.

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Do Not Breathe Dust. Do Not Take Internally. Avoid Breathing of Dust. Skin Contact May Be Harmful. Avoid Contact With Skin. Wash Hands Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily. Food utensils such as tablespoons should not be used for food purposes after use with pesticides. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protection information may be obtained from your Cooperative Agricultural Extension Service. Do not reuse empty container. Destroy it by burying waste or burning. Stay away from smoke or fumes. Note to Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS.**

DIRECTIONS FOR USE: BLACK LEAF TOMATO AND VEGETABLE DUST may be used in the conventional plants, spray dust or also as a liquid spray. Best Results will be obtained when spraying in early morning or late evening during periods of low temperature and a movement. **BLACK LEAF TOMATO AND VEGETABLE DUST** should be applied to all exposed parts of the plant including the lower leaf surfaces.


SPRAYING: Mix three ounces (1 1/2 tablespoons) in about 1 pint of water until all of the dust is moist or partly so, then add to 1 gallon of water. During the spraying, please continue to shake or agitate. Avoid burning this diluted spray.

BLACK LEAF TOMATO AND VEGETABLE DIRT should be used only as a bulk form to apply. The proper dilution should be reported at 5 to 10 day intervals.

FUNGUS CONTROL: To control the many diseases that can afflict your grapevines, protect them from fungi. Buy fungicides labeled for use on grapes. Apply fungicides to the leaves, stems, and fruit. Do this before and after rain, and after a frost, and before and after a hail storm.

NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Contains 5% SEVIN-6% MANEB



Controls insects and diseases of plants and animals

A double action easy-to-use garden dust

**CAUTION:
KEEP OUT OF THE
REACH OF CHILDREN.**

NET WEIGHT 100g

BLACK LEAF PRODUCTS COMPANY

100-111013-60120

Contains 5% SEVIN-6% MANEB

Controls insects and diseases on tomatoes as well as other vegetables

**A double action
easy-to-use
garden dust**

**CAUTION:
KEEP OUT OF THE
REACH OF CHILDREN**

NET WEIGHT 10 OZ

E.P.A. Reg. No. 5887-40

ACTIVE INGREDIENTS

SEVIN (Fenitrothion)	5.0%
MANEB (Manganese Ethylenebisdithiocarbamate)	6.0%
INERT INGREDIENTS	89.0%

BLACK LEAF PRODUCTS COMPANY

Chicago, Illinois 60620

lower leaf surfaces.

SPRAYING: Mix three ounces (11 tabs) about 1 pint of water until all of the dust is dissolved or pasty and then add to 1 gallon of water. During the spraying please continue to shake the container. Avoid stirring this diluted spray.

BLACK LEAF TOMATO AND VEGETABLE DUST should be used as soon as buds begin to open. The spray or dusting should be repeated at 7 to 10 day intervals.

FUNGUS CONTROL: To control the mildew and blight that can affect your vegetables a dusting of **BLACK LEAF TOMATO AND VEGETABLE DUST** should commence as soon as foliage appears and repeat every 1 or 2 weeks.

NOTICE: Buyer assumes all risks of injury or damage from the use or handling of this material not in strict accordance with directions given herewith.